

Renormalization group invariants in the MSSM and its extensions

This content has been downloaded from IOPscience. Please scroll down to see the full text.

JHEP11(2005)003

(<http://iopscience.iop.org/1126-6708/2005/11/003>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 193.140.249.2

This content was downloaded on 18/07/2016 at 09:14

Please note that [terms and conditions apply](#).

Renormalization group invariants in the MSSM and its extensions

Durmuş A. Demir

*Department of Physics, Izmir Institute of Technology, IZTECH
Turkey, TR35430
E-mail: demir@physics.iztech.edu.tr*

ABSTRACT: We derive one-loop renormalization group (RG) invariant observables and analyze their phenomenological implications in the MSSM and its μ problem solving extensions, $U(1)'$ model and NMSSM. We show that there exist several RG invariants in the gauge, Yukawa and soft-breaking sectors of each model. In general, RG invariants are highly useful for projecting experimental data to messenger scale, for revealing correlations among the model parameters, and for probing the mechanism that breaks supersymmetry. The Yukawa couplings and trilinear soft terms in $U(1)'$ model and NMSSM do not form RG invariants though there exist approximate invariants in low $\tan\beta$ domain. In the NMSSM, there are no invariants that contain the Higgs mass-squareds. We provide a comparative analysis of RG invariants in all three models and analyze their model-building and phenomenological implications by a number of case studies.

KEYWORDS: Renormalization Group, Supersymmetry Phenomenology.

Contents

1. Introduction	1
2. Generalities	3
3. The RG invariants in the MSSM	4
4. The RG invariants in $U(1)'$ Model	11
5. The RG invariants in the NMSSM	17
6. Conclusion	20
A. Renormalization group equations in $U(1)'$ model	22
B. Renormalization group equations in the NMSSM	25

1. Introduction

The supersymmetric models provide an elegant solution to the gauge hierarchy problem in a genuinely perturbative way for all scales right up to the planckian territory. The perturbative nature of the model allows one to relate measurements at the electroweak scale to physics at ultra high energies. This communication between the infrared (IR) and ultra-violet (UV) regimes proceeds with the renormalization group (RG) flow of the lagrangian parameters. Indeed, various phenomena central to supersymmetry phenomenology e.g. gauge coupling unification [1], radiative electroweak breaking [2], induction of flavor structures [3] even for flavor-blind soft terms are pure renormalization effects.

Projection of experimental data to ultra high energies requires solving renormalization group equations (RGEs) for parameters of the model. This projection, however, is generally complicated by the coupled nature of RGEs in that measurement of a set of parameters at low scale cannot directly be rescaled to ultra high energies due to leakage of other, possibly unknown, quantities. Therefore, in course of fitting a given model to laboratory and astrophysical data it would be advantageous to have as much information as possible about correlations among the parameters. Concerning this, a highly useful tool is provided by the RG-invariant observables. Indeed, such quantities prove highly useful not only for projecting the experimental data to high energies but also for deriving certain sum rules which enable fast consistency checks of the model [4–6]. However, it should be kept in mind that, even the RG-invariant observables cannot be guaranteed to work perfectly because (i) the RG invariance holds at a given loop order and it is generically disrupted

by higher loop effects (In general, scale and conformal invariances imply each other [7], and superconformal group involves both scale invariance and a continuous R symmetry with correlated charges [8]. Therefore, in softly broken supersymmetric theories, where R invariance is explicitly broken, the RG invariance, if any, cannot be an all-order effect.), (ii) the RGEs get modified at sparticle thresholds so that what equations must be used is not known *a priori*, and (iii) parameters with nontrivial flavor structures typically do not exhibit RG-invariant combinations (this can be tied up to the fact that scale invariance puts strong constraints on flavor structures of rigid and soft parameters of the theory [9]). The flavor mixings and sparticle thresholds disrupt RG equations and associated invariants already at one loop order. On the other hand, modification in a given RG invariant due to higher loop contributions is of order one less loop factor. For instance, disruption of a one-loop RG invariant by two loop effects is of one loop order. Below we will restrict our analysis to one loop RGEs with no flavor mixings in fermion sector.

This work is devoted to derivations and analyses of RG-invariant observables in the minimal supersymmetric model (MSSM) and its μ -problem solving minimal extensions i.e. the next-to-minimal supersymmetry (NMSSM) and its gauged version $U(1)'$ model. Indeed, MSSM suffers from the naturalness problem associated with the Dirac mass of Higgsinos. This mass parameter, μ , is nested in the superpotential of the theory, and hence, its scale is not controlled by the mechanism that breaks supersymmetry [10]. Consequently, it is necessary to find a mechanism for stabilizing μ to the electroweak scale. In fact, $U(1)'$ model and NMSSM both provide a dynamical solution to the problem by inducing μ via the VEV of an MSSM-singlet chiral superfield. The $U(1)'$ models are extensions of the MSSM by both an MSSM singlet and an additional abelian symmetry $U(1)'$ [11, 12]. On the other hand, NMSSM has the same gauge group as MSSM yet its spectrum contains a pure gauge singlet [13]. One here notes that a TeV scale $U(1)'$ symmetry or NMSSM are not necessarily the only solutions to the μ problem. Indeed, modification of the Kahler potential by operators of the form $M_{Pl}^{-1} \hat{z}^\dagger \hat{H}_u \hat{H}_d$, \hat{z} being a hidden sector field, generates the μ parameter at the right scale provided that theory possesses a global Peccei-Quinn or continuous R invariance to forbid a bare μ parameter to appear in the superpotential [14] (see also the related scenarios in [15] and [16]). Apart from this, the mechanism proposed in [17] provides a simultaneous solution to the μ problem and the scale of supersymmetry breaking within the supergravity framework by constructing explicit models of the hidden sector.

The RG-invariant observables and their phenomenological implications have already been analyzed in various contexts. In addition to discussions in [4–6] there have been studies of the RG invariants [18] and resulting sum rules [19] within supersymmetric gauge theories and certain string-inspired soft terms. In this work, we will provide a comparative analysis of the RG invariants in the MSSM and its μ problem solving extensions. The RGEs for $U(1)'$ model had been first given in [12]. Here we generalize them to finite bottom and tau Yukawas. They are listed in appendix A. The RGEs for NMSSM had been given in [20], and we rederive and list them in appendix B, for completeness. In appendices we also discuss limiting cases where $U(1)'$ and NMSSM RGEs reduce to those of the MSSM [21].

We start our analysis, in section 2 below, by first describing the common part of all three models i.e. the sfermion sector. Then we derive RG-invariant observables and discuss their phenomenological implications for the MSSM in section 3, for $U(1)'$ model in section 4, and finally for the NMSSM in section 5. In section 6 we conclude the work.

2. Generalities

For the three supersymmetric models we will discuss the fermion sector is common whereas the Higgs and gauge sectors vary from model to model. In general, one can write

$$\widehat{W} = \widehat{W}_{\text{Higgs}} + \widehat{W}_{\text{fermion}} \quad (2.1)$$

where the superpotential of the fermion sector is given by

$$\widehat{W}_{\text{fermion}} = \widehat{U} \mathbf{Y}_{\mathbf{u}} \widehat{Q} \widehat{H}_u + \widehat{D} \mathbf{Y}_{\mathbf{d}} \widehat{Q} \widehat{H}_d + \widehat{E} \mathbf{Y}_{\mathbf{e}} \widehat{L} \widehat{H}_d \quad (2.2)$$

which encodes the Yukawa couplings $\mathbf{Y}_{\mathbf{u,d,e}}$ (of up quarks, down quarks and of leptons) each being a 3×3 non-hermitean matrix in the space of fermion flavors. The fermion masses are induced by the vacuum expectation values of the Higgs doublets \widehat{H}_u and \widehat{H}_d , common to all three models. In $\widehat{W}_{\text{fermion}}$ the left-handed quarks are arranged in $SU(2)_L$ doublets \widehat{Q} and the left-handed leptons in $SU(2)_L$ doublet \widehat{L} . On the other hand, the left-handed anti up and anti down quarks are represented by \widehat{U} and \widehat{D} , respectively. Finally, \widehat{E} collects left-handed anti leptons.¹

The breakdown of supersymmetry is parameterized by various soft terms belonging to Higgs, gaugino and scalar fermion sectors (see, e.g. the review volume [22]):

$$\mathcal{L}_{\text{soft}} = \mathcal{L}_{\text{soft}}^{\text{Higgs}} + \mathcal{L}_{\text{soft}}^{\text{gaugino}} + \mathcal{L}_{\text{soft}}^{\text{sfermion}} \quad (2.3)$$

whose sfermion part reads as

$$-\mathcal{L}_{\text{soft}}^{\text{sfermion}} = \widetilde{Q}^\dagger \mathbf{m}_{\mathbf{Q}}^2 \widetilde{Q} + \widetilde{U} \mathbf{m}_{\mathbf{U}}^2 \widetilde{U}^\dagger + \widetilde{D} \mathbf{m}_{\mathbf{D}}^2 \widetilde{D}^\dagger + \widetilde{L}^\dagger \mathbf{m}_{\mathbf{L}}^2 \widetilde{L} + \widetilde{E} \mathbf{m}_{\mathbf{E}}^2 \widetilde{E}^\dagger + \left[\widetilde{U} \mathbf{Y}_{\mathbf{u}}^{\mathbf{A}} \widetilde{Q} \widehat{H}_u + \widetilde{D} \mathbf{Y}_{\mathbf{d}}^{\mathbf{A}} \widetilde{Q} \widehat{H}_d + \widetilde{E} \mathbf{Y}_{\mathbf{e}}^{\mathbf{A}} \widetilde{L} \widehat{H}_d + \text{h.c.} \right], \quad (2.4)$$

where $\mathbf{Y}_{\mathbf{u,d,e}}^{\mathbf{A}}$, like Yukawas themselves, are non-hermitean flavor matrices whereas the sfermion mass-squareds $\mathbf{m}_{\mathbf{Q},\dots,\mathbf{E}}^2$ are all hermitean.

The interactions contained in (2.2) and (2.4) exhibit mixings of various flavors in both rigid and soft terms. As mentioned in Introduction, we focus only on the flavor-diagonal interactions due to the fact that flavor mixings generically prohibit the construction of RG invariants except for those parameters which depend on traces or determinants of the flavor matrices. Moreover, one recalls that there is a certain degree of correlation between conformal invariance and flavor violation in that the former can put stringent constraints on the latter [9]. Consequently, we switch off flavor mixings in all rigid and soft parameters

¹The neutrino masses and hence the requisite superfields (heavy Majorana neutrinos or light right-handed neutrinos) are not incorporated in supersymmetric models under discussion.

to obtain

$$\begin{aligned}
 \mathbf{Y}_{\mathbf{u,d,e}} &\rightarrow \text{diag. } (0, 0, h_{t,b,\tau}), \\
 \mathbf{Y}_{\mathbf{u,d,e}}^{\mathbf{A}} &\rightarrow \text{diag. } (0, 0, h_{t,b,\tau} A_{t,b,\tau}), \\
 \mathbf{m}_{\mathbf{Q}}^2 &\rightarrow \text{diag. } (m_{\tilde{u}_L}^2, m_{\tilde{c}_L}^2, m_{\tilde{t}_L}^2), \\
 \mathbf{m}_{\mathbf{L}}^2 &\rightarrow \text{diag. } (m_{\tilde{e}_L}^2, m_{\tilde{\mu}_L}^2, m_{\tilde{\tau}_L}^2), \\
 \mathbf{m}_{\mathbf{U}}^2 &\rightarrow \text{diag. } (m_{\tilde{u}_R}^2, m_{\tilde{c}_R}^2, m_{\tilde{t}_R}^2), \\
 \mathbf{m}_{\mathbf{D}}^2 &\rightarrow \text{diag. } (m_{\tilde{d}_R}^2, m_{\tilde{s}_R}^2, m_{\tilde{b}_R}^2), \\
 \mathbf{m}_{\mathbf{E}}^2 &\rightarrow \text{diag. } (m_{\tilde{e}_R}^2, m_{\tilde{\mu}_R}^2, m_{\tilde{\tau}_R}^2),
 \end{aligned} \tag{2.5}$$

where $m_{\tilde{u}_L}^2 = m_{\tilde{d}_L}^2$, $m_{\tilde{c}_L}^2 = m_{\tilde{s}_L}^2$ and $m_{\tilde{t}_L}^2 = m_{\tilde{b}_L}^2$ by gauge invariance. Note that light fermion Yukawa couplings are totally neglected. This reduction scheme for flavor mixings sets up the notation and framework for the fermion sector. The gauge and Higgs sectors differ from model to model, and they will be discussed in detail in the following sections.

Another model-independent aspect to be noted concerns IR and UV boundaries of the RGEs. For all three supersymmetric models of interest, we neglect modifications in the particle spectrum and RGEs coming from decoupling of the heavy fields. In other words, we assume that all soft masses are approximately equal to $M_{\text{SUSY}} \sim 1 \text{ TeV}$ in logarithmic sense. This scale sets up the IR boundary for exact supersymmetric RG flow. The UV boundary lies just beneath the scale of string territory, and we will take it to be the scale of gauge coupling unification in the MSSM: $M_{\text{GUT}} \sim 10^{16} \text{ GeV}$. Therefore, in our framework, the RG invariance of a given quantity means its scale independence in between the IR and UV scales above. In what follows, we judiciously combine the RGEs of individual quantities until we arrive at a RG-invariant observable within one loop accuracy. In general, there is no guarantee of maintaining RG invariance of a given quantity at higher loop levels.

3. The RG invariants in the MSSM

The MSSM is based on $\text{SU}(3)_c \times \text{SU}(2)_L \times \text{U}(1)_Y$ gauge group with respective gauge couplings g_3 , g_2 and g_1 . The Higgs sector is spanned by \widehat{H}_u and \widehat{H}_d so that

$$\widehat{W}_{\text{Higgs}} = \mu \widehat{H}_u \widehat{H}_d \tag{3.1}$$

and

$$\begin{aligned}
 -\mathcal{L}_{\text{soft}}^{\text{Higgs}} &= m_{H_u}^2 H_u^\dagger H_u + m_{H_d}^2 H_d^\dagger H_d + [\mu B H_u H_d + \text{h.c.}], \\
 -\mathcal{L}_{\text{soft}}^{\text{gaugino}} &= \frac{1}{2} \sum_{a=3,2,1} [M_a \lambda_a \lambda_a + \text{h.c.}],
 \end{aligned} \tag{3.2}$$

where M_a is the gaugino mass.

By using one-loop RGEs (the RGEs in the MSSM have been computed up to two (partially up to three) loops order in [21]) within the MSSM one can derive a number of invariants. Several RG invariants are listed in table 1. The invariant I_1 correlates the gauge couplings with arbitrary constants c_1 and c_2 . The constants, however, can be related by using the values of gauge couplings at M_{SUSY} and M_{GUT} :

$$\frac{c_1}{c_2} = -\frac{5 g_1(M_{\text{SUSY}})^2}{3 g_2(M_{\text{SUSY}})^2} \left(\frac{g_0^2 g_2(M_{\text{SUSY}})^2 + 3 g_0^2 g_3(M_{\text{SUSY}})^2 - 4 g_2(M_{\text{SUSY}})^2 g_3(M_{\text{SUSY}})^2}{11 g_0^2 g_1(M_{\text{SUSY}})^2 + 5 g_0^2 g_3(M_{\text{SUSY}})^2 - 16 g_1(M_{\text{SUSY}})^2 g_3(M_{\text{SUSY}})^2} \right), \quad (3.3)$$

where g_0 is the common value of the gauge couplings at the unification scale M_{GUT} .

The second invariant I_2 in table 1 correlates μ parameter with gauge and Yukawa couplings. From this one can determine μ at any scale $Q \in [M_{\text{SUSY}}, M_{\text{GUT}}]$:

$$\begin{aligned} \mu(Q_2) = & \mu(Q_1) \left(\frac{h_t(Q_2)}{h_t(Q_1)} \right)^{27/61} \left(\frac{h_b(Q_2)}{h_b(Q_1)} \right)^{21/61} \left(\frac{h_\tau(Q_2)}{h_\tau(Q_1)} \right)^{10/61} \times \\ & \times \left(\frac{g_3(Q_1)}{g_3(Q_2)} \right)^{256/183} \left(\frac{g_2(Q_1)}{g_2(Q_2)} \right)^{9/61} \left(\frac{g_1(Q_2)}{g_1(Q_1)} \right)^{73/2013} \end{aligned} \quad (3.4)$$

which makes it manifest that μ at any scale Q depends on the strong coupling g_3 although its RGE does not exhibit such a direct dependence at all. This exemplifies one interesting aspect of the RG invariants: they make various otherwise implicit dependencies explicit. By putting $Q_2 = M_{\text{SUSY}}$ and $Q_1 = M_{\text{GUT}}$ one finds that the ratio $\mu(M_{\text{SUSY}})/\mu(M_{\text{GUT}})$, which is one of the most crucial factors (together with the gluino mass) that determine the amount of fine-tuning needed to achieve the correct value of the Z boson mass, is entirely determined by the interplay between the IR and UV values of the rigid parameters. In particular, (3.4) suggests that $\mu(M_{\text{SUSY}})/\mu(M_{\text{GUT}})$ decreases with increasing $\tan\beta$: $\mu(M_{\text{SUSY}})/\mu(M_{\text{GUT}}) \simeq 0.96$ for $\tan\beta = 5$ and $\simeq 0.3$ for $\tan\beta = 60$. Indeed, this ratio is governed mainly by g_3 at low $\tan\beta$ and by g_3 , h_b and h_τ for $\tan\beta \sim m_t/m_b$. Therefore, the sensitivity of M_Z to $\mu(M_{\text{GUT}})$ is greatly reduced at large $\tan\beta$ which itself requires a great deal of fine-tuning to achieve though [23] (see [24] for a discussion of the fine-tuning problems in large $\tan\beta$ domain when radiative corrections to Higgs potential are taken into account).

The third line of table 1 shows that the ratio of the gaugino mass to fine structure constant of the same group is an RG invariant. This invariance property guarantees that

$$M_a(Q_2) = M_a(Q_1) \left(\frac{g_a(Q_2)}{g_a(Q_1)} \right)^2 \quad (3.5)$$

so that knowing two of the gaugino masses at a scale Q suffices to know the third if gauge coupling unification holds — an important aspect to check directly the minimality of the gauge structure using the experimental data. This very relation also shows that $M_3(M_{\text{SUSY}})/M_3(M_{\text{GUT}})$ is much larger $M_{1,2}(M_{\text{SUSY}})/M_{1,2}(M_{\text{GUT}})$ due to asymptotic freedom. In fact, in minimal supergravity for instance, typically gluino is the first superpartner to decouple from the light spectrum.

Number	RG Invariant
I_1	$\frac{c_1}{g_1^2} + \frac{c_2}{g_2^2} + \frac{33c_1+5c_2}{15g_3^2}$
I_2	$\mu \left(\frac{g_2^9 g_3^{256/3}}{h_t^{27} h_b^{21} h_\tau^{10} g_1^{73/33}} \right)^{1/61}$
I_3	$\frac{M_a}{g_a^2} \quad (a = 1, 2, 3)$
I_4	$B - \frac{27}{61}A_t - \frac{21}{61}A_b - \frac{10}{61}A_\tau - \frac{256}{183}M_3 - \frac{9}{61}M_2 + \frac{73}{2013}M_1$
I_5	$m_{\tilde{\tau}_R}^2 - 2m_{\tilde{\tau}_L}^2 - 3 M_2 ^2 + \frac{1}{11} M_1 ^2 - \frac{6}{13}S$
I_6	$m_{H_u}^2 - \frac{3}{2}m_{\tilde{t}_R}^2 + \frac{4}{3} M_3 ^2 + \frac{3}{2} M_2 ^2 - \frac{5}{66} M_1 ^2 - \frac{9}{26}S$
I_7	$m_{H_d}^2 - \frac{3}{2}m_{\tilde{b}_R}^2 - m_{\tilde{\tau}_L}^2 + \frac{4}{3} M_3 ^2 - \frac{1}{33} M_1 ^2 + \frac{3}{26}S$
I_8	$m_{\tilde{t}_R}^2 + m_{\tilde{b}_R}^2 - 2m_{\tilde{t}_L}^2 - 3 M_2 ^2 + \frac{1}{11} M_1 ^2 + \frac{2}{13}S$
I_9	$m_{\tilde{u}_L}^2 + \frac{1}{198} M_1 ^2 + \frac{3}{2} M_2 ^2 - \frac{8}{9} M_3 ^2 - \frac{1}{26}S$
I_{10}	$m_{\tilde{u}_R}^2 + \frac{8}{99} M_1 ^2 - \frac{8}{9} M_3 ^2 + \frac{2}{13}S$
I_{11}	$m_{\tilde{d}_R}^2 + \frac{2}{99} M_1 ^2 - \frac{8}{9} M_3 ^2 - \frac{1}{13}S$
I_{12}	$m_{\tilde{e}_L}^2 + \frac{1}{22} M_1 ^2 - \frac{3}{2} M_2 ^2 + \frac{3}{26}S$
I_{13}	$m_{\tilde{e}_R}^2 + \frac{2}{11} M_1 ^2 - \frac{3}{13}S$
I_{14}	$m_{H_u}^2 + m_{H_d}^2 - 3m_{\tilde{t}_L}^2 - m_{\tilde{\tau}_L}^2 + \frac{8}{3} M_3 ^2 - 3 M_2 ^2 + \frac{2}{11} M_1 ^2$
I_{15}	$m_{H_d}^2 - \frac{3}{2}m_{\tilde{b}_R}^2 - \frac{3}{2}m_{\tilde{\tau}_L}^2 + \frac{1}{4}m_{\tilde{\tau}_R}^2 + \frac{4}{3} M_3 ^2 - \frac{3}{4} M_2 ^2 - \frac{1}{132} M_1 ^2$
I_{16}	$2m_{\tilde{u}_L}^2 + m_{\tilde{u}_R}^2 + m_{\tilde{d}_R}^2 + \frac{1}{9} M_1 ^2 + 3 M_2 ^2 - \frac{32}{9} M_3 ^2$
I_{17}	$m_{\tilde{e}_L}^2 + \frac{1}{2}m_{\tilde{e}_R}^2 + \frac{3}{22} M_1 ^2 - \frac{3}{2} M_2 ^2$

Table 1: The RG invariant combinations of rigid and soft parameters in the MSSM (c_1 and c_2 in I_1 are arbitrary constants). Note that invariants pertaining to the first and second generations generically involve a single sfermion mass-squared since trilinear couplings do not contribute to their RGEs. In a sense, these are 'fundamental' invariants derived directly from the RG flows of relevant parameters. For obtaining RG invariants containing a specific set of parameters it is necessary to form appropriate combinations of these tabulated ones, as exemplified in the text by a couple of case studies. The quantity S appearing in some of the invariants is defined in equation (3.7).

The fourth line of table 1 correlates Higgs bilinear soft term B with trilinear couplings and gaugino masses. Among various possibilities, by using this invariant one can express, for instance, B at any scale Q in terms of other dimension-one soft masses:

$$\begin{aligned}
 B(Q_2) = & B(Q_1) + \frac{27}{61} (A_t(Q_2) - A_t(Q_1)) + \frac{21}{61} (A_b(Q_2) - A_b(Q_1)) + \\
 & + \frac{10}{61} (A_\tau(Q_2) - A_\tau(Q_1)) + \frac{256}{183} M_3(Q_1) \left(\frac{g_3(Q_2)^2}{g_3(Q_1)^2} - 1 \right) + \\
 & + \frac{9}{61} M_2(Q_1) \left(\frac{g_2(Q_2)^2}{g_2(Q_1)^2} - 1 \right) - \frac{73}{2013} M_1(Q_1) \left(\frac{g_1(Q_2)^2}{g_1(Q_1)^2} - 1 \right) \quad (3.6)
 \end{aligned}$$

after using (3.5). This equation expresses the IR value of the B parameter in terms of the IR and UV values of the gaugino masses and trilinear couplings. The RGE of the B parameter does not depend on the gluino mass explicitly (the dependence comes through the trilinear couplings); however, (3.6) exhibits a rather strong dependence on M_3 : for

$Q_2 = M_{\text{SUSY}}$ and $Q_1 = M_{\text{GUT}}$ the gluino contribution equals $2.6M_3(M_{\text{GUT}})$ which is much larger than other contributions (except possibly the GUT scale value of B). This very fact proves the power of forming RG invariant observables as they make indirect effects manifest.

Having completed the discussion of the rigid and dimension-one soft parameters of the theory, we now start analyzing the scale-invariant combinations of the scalar mass-squareds. They are listed in table 1 starting from line 5. The RGEs of the soft mass-squareds depend on the quantity [21]

$$\begin{aligned}
 S = \text{Tr} [m^2 Y] = & m_{H_u}^2 - m_{H_d}^2 + (m_{t_L}^2 - m_{t_R}^2) + (m_{c_L}^2 - m_{c_R}^2) + (m_{u_L}^2 - m_{u_R}^2) + \\
 & + (m_{b_R}^2 - m_{t_R}^2) + (m_{s_R}^2 - m_{c_R}^2) + (m_{d_R}^2 - m_{u_R}^2) - \\
 & - (m_{\tau_L}^2 - m_{\tau_R}^2) - (m_{\mu_L}^2 - m_{\mu_R}^2) - (m_{e_L}^2 - m_{e_R}^2)
 \end{aligned}
 \tag{3.7}$$

which comprises all of the soft mass-squareds. This quantity identically vanishes if they are strictly universal at some given scale since then $\text{Tr}[m^2 Y] = m^2 \text{Tr}[Y] \equiv 0$ thanks to the absence of the gravitational anomaly. As the explicit solution

$$S(Q_2) = \left(\frac{g_1(Q_2)}{g_1(Q_1)} \right)^{26/33} S(Q_1)
 \tag{3.8}$$

also suggests, $S(Q)$ vanishes at all scales if it does so at some given scale. That the universality of the soft mass-squareds,

$$m_{H_u}^2 = m_{H_d}^2 = \dots = m_{\tau_R}^2 = m_0^2,
 \tag{3.9}$$

renders theory S -free is important in that experimental tests of whether S is vanishing or not can give important information on if soft masses unify at ultra high energies. This universality scheme, when supplemented by $A_t(M_{\text{GUT}}) = A_b(M_{\text{GUT}}) = A_\tau(M_{\text{GUT}}) = A_0$ and $M_3(M_{\text{GUT}}) = M_2(M_{\text{GUT}}) = M_1(M_{\text{GUT}}) = M$, leads one to the minimal supergravity configuration. One further notes that, the RGE of a scalar ϕ senses S via the contribution $Y_\phi(3/5)g_1^2 S$; however, the RG invariants of the soft mass-squareds depend on S without any g_1^2 dressing.

The RG-invariant combinations of the soft mass-squareds can therefore be analyzed in two groups: those that are sensitive to S (lines 5-13 of table 1) and those that are insensitive to S (lines 14-17 of table 1). Clearly, one can construct new invariants by combining these available ones. An accurate enough measurement (presumably at $\text{LHC} \oplus \text{ILC}$) of (all or part of the) soft mass-squareds will serve both as a testing ground for the internal consistency of the model and as a tool for probing the ultra high energy behavior (whether it is minimal supergravity or not within experimental error bounds) [19, 25]. More specifically, by using these invariants one can (i) test the internal consistency of the model while fitting to the experimental data; (ii) rehabilitate poorly known parameters supplementing the well-measured ones; (iii) determine what kind of supersymmetry breaking mechanism is realized in Nature; and finally (iv) separately examine the UV scale configurations of the

trilinear couplings as they do not explicitly contribute to these invariants. We now want to illustrate some of these useful aspects of these invariants by studying a few interesting cases:

- The RG invariants provide useful constraints on the low-energy parameter space. For illustrating this point let us consider, for example, the RG invariant I_8 in table 1. If the universality constraint (3.9) holds at some scale Q_1 then one finds

$$m_{\tilde{t}_R}^2(Q_2) + m_{\tilde{b}_R}^2(Q_2) - 2m_{\tilde{t}_L}^2(Q_2) = 3|M_2(Q_1)|^2 \left[\left(\frac{g_2(Q_2)}{g_2(Q_1)} \right)^4 - 1 \right] - \frac{1}{11} |M_1(Q_1)|^2 \left[\left(\frac{g_1(Q_2)}{g_1(Q_1)} \right)^4 - 1 \right] \quad (3.10)$$

after using equation (3.5). This equality establishes a relation between the stop and sbottom masses right at the scale of measurement in a way involving the UV values of the gaugino masses. Despite this, however, it shows that $\tilde{t}_L - \tilde{t}_R$ plus $\tilde{b}_L - \tilde{b}_R$ mass splittings are entirely controlled by the isospin and hypercharge gaugino masses rather than the gluino mass. This is an important aspect as it significantly reduces sensitivity to the UV scale values of the gaugino masses. Indeed, by taking $Q_2 = M_{\text{SUSY}}$ and $Q_1 = M_{\text{GUT}}$ the right-hand side of (3.10) reduces to $-0.97M_2(M_{\text{GUT}})^2 + 0.08M_1(M_{\text{GUT}})^2$ which does not exhibit any pronounced sensitivity to GUT scale gaugino masses (unlike, for instance, (3.6)). One possible application of (3.10) among many one can consider is that it establishes a relation between the stop and sbottom mixing angles

$$\sum_{f=t,b} \frac{m_f |A_f - \mu^* R_f|}{\tan 2\theta_{\tilde{f}}} \simeq -0.5M_2(M_{\text{GUT}})^2 + 0.04M_1(M_{\text{GUT}})^2 - 0.04M_2^2, \quad (3.11)$$

where $R_f = \cot \beta (\tan \beta)$ for $f = t(b)$. In estimating the right hand side we took $\cos 2\beta \simeq -1$ in accord with the LEP bounds which prefer fairly large values of $\tan \beta$. This simple formula may serve as a constraint in simulating the supersymmetric parameter space as the experimental data accumulate. So far we have assumed that the theory is S -free. What if it is not? In this case one automatically obtains a direct sensitivity to all soft mass-squareds and neither (3.10) nor (3.11) can provide a signal as clean as in the universal case.

- The RG invariants in table 1 can be combined to obtain new invariants that involve solely the scalar mass-squareds in the theory. For instance, by taking M_{GUT} to be the UV scale with universal scalar masses the Higgs soft masses can be expressed as

$$\begin{aligned} m_{H_u}^2(Q) &= \frac{7}{12}m_0^2 + \frac{5}{12}m_{\tilde{e}_R}^2(Q) + m_{\tilde{u}_L}^2(Q) - \frac{21}{12}m_{\tilde{u}_R}^2(Q) - \frac{3}{4}m_{\tilde{d}_R}^2(Q) + \frac{3}{2}m_{\tilde{t}_R}^2(Q) \\ m_{H_d}^2(Q) &= -\frac{15}{4}m_0^2 - \frac{1}{4}m_{\tilde{e}_R}^2(Q) - 3 \left(m_{\tilde{u}_L}^2(Q) - \frac{1}{2}m_{\tilde{u}_R}^2(Q) \right) - \\ &\quad - 3 \left(m_{\tilde{t}_L}^2(Q) - \frac{1}{2}m_{\tilde{t}_R}^2(Q) - \frac{3}{4}m_{\tilde{b}_R}^2(Q) \right) + \frac{3}{2} \left(m_{\tilde{\tau}_L}^2(Q) - \frac{1}{6}m_{\tilde{\tau}_R}^2(Q) \right) \end{aligned} \quad (3.12)$$

which serve as a sum rule to correlate various soft masses with no contributions from those of the gauginos. These expressions determine Higgs soft mass-squareds in terms of the sfermion masses and the universal scalar mass at the GUT scale. Clearly, for determining the Higgs soft mass-squareds it is necessary to know soft masses of all three generations of sfermions if the gaugino sector is to be skipped (compare (3.12) with the invariants I_6 , I_7 and I_{14} in table 1).

The RG invariant I_{13} can prove useful for determining S from a minimal number of measurements. Indeed, measuring $m_{\tilde{e}_R}$ and M_1 at two distinct scales and taking the difference determines if S is vanishing (if scalar masses attain a universal scheme as in (3.9)) or not. However, given that a measurement of M_1 requires exploration of the neutralino sector, a more promising invariant is

$$m_{\tilde{u}_R}^2 - m_{\tilde{d}_R}^2 - \frac{1}{3}m_{\tilde{e}_R}^2 + \frac{4}{13}S \tag{3.13}$$

which involves only the first family sfermions which are simultaneous eigenstates of mass, gauge, flavor and chirality (and thus, their experimental identification could be easier than those of gauginos and third generation sfermions which undergo non-negligible mixings).

- The RG invariants are highly useful probes of the mechanism that breaks the supersymmetry. We illustrate their discriminative power by examining three well-known supersymmetry breaking schemes: (i) no-scale supergravity models [26], (ii) dilaton-dominated supersymmetry breaking [27] and (iii) flux-induced soft terms [28]. The soft-breaking sectors of these models commonly exhibit the minimal supergravity (constrained MSSM) configuration: $m_{H_u}^2 = m_{H_d}^2 = \dots = m_{\tilde{\tau}_R}^2 = m_0^2$, $A_t(M_{\text{GUT}}) = A_b(M_{\text{GUT}}) = A_\tau(M_{\text{GUT}}) = A_0$ and $M_3(M_{\text{GUT}}) = M_2(M_{\text{GUT}}) = M_1(M_{\text{GUT}}) = M$. However, correlations among the parameters vary from model to model so does the pattern of the RG invariants. In no-scale supergravity $A_0 = B(M_{\text{GUT}}) = m_0 = 0$, in dilaton domination $m_0 = M/\sqrt{3}$, $A_0 = -M$ and $B(M_{\text{GUT}}) = 2M/\sqrt{3}$, and in fluxed MSSM $m_0 = M$, $A_0 = -3M$ and $B(M_{\text{GUT}}) = -2M$. The values of the soft-sector invariants are displayed in tables 2 and 3. One notices that, the only model-independent invariant is I_8 as it solely probes if soft mass-squareds and/or gaugino masses are universal or not. The other invariants differ from model to model. In case one invariant, say I_4 , is determined by experiment with sufficient accuracy and if it agrees with predictions of a specific model, say dilaton-dominated supersymmetry breaking, then all one has to do is to check if rest of the invariants (to be determined as more and more data accumulate) agree with the experiment. In this sense, the results displayed in tables 2 and 3 (which can be expanded to include all possible breaking schemes found in strings, supergravity, anomaly mediation, gauge mediation, etc.) can be used as a look up table for checking/predicting which mechanism of supersymmetry breaking is favored or realized in Nature. Clearly, RG invariance is not a requisite property for an observable to probe supersymmetry breaking sector; however, if it is RG-invariant it is not necessary to integrate the RGEs and it is possible to use results

Mechanism	I_4/M	I_5/M^2	I_6/M^2	I_7/M^2	I_8/M^2	I_9/M^2
No-scale	-3040/2013	-32/11	91/33	43/33	-32/11	61/99
Dilaton	$2/\sqrt{3}-1126/2013$	-107/33	57/22	53/66	-32/11	94/99
Fluxed	-7462/2013	-43/11	149/66	-13/66	-32/11	160/99

Table 2: The patterns of the RG invariants (I_4 - I_9) within no-scale [26], dilaton domination [27] and fluxed MSSM [28] supersymmetry breaking schemes.

Mechanism	I_{10}/M^2	I_{11}/M^2	I_{12}/M^2	I_{13}/M^2	I_{14}/M^2	I_{15}/M^2	I_{16}/M^2	I_{17}/M^2
No-scale	-80/99	-86/99	-16/11	2/11	-10/33	19/33	-4/9	-15/11
Dilaton	-47/99	-53/99	-37/33	17/33	-32/33	-1/132	8/9	-19/22
Fluxed	19/99	13/99	-5/11	13/11	-76/33	-155/132	32/9	3/22

Table 3: The same as in table 2 but for I_{10} - I_{17} .

of different experiments without RG scaling (irrespective of if they are obtained from a reanalysis of LEP data or from Tevatron or from LHC).

- The RG invariants do have interesting implications also for chargino and neutralino sectors of the model. For instance, from the product of two chargino masses one immediately finds that the quantity

$$\left(M_{\chi_1^\pm} M_{\chi_2^\pm} + M_W^2 \sin 2\beta \right) \left(\frac{g_3^{256/3}}{h_t^{27} h_b^{21} h_\tau^{10} g_2^{113} g_1^{73/33}} \right)^{1/61} \quad (3.14)$$

is an RG invariant observable of the model. Besides, sum of the neutralino masses can be shown to be expressible in terms of the gauge couplings and gluino mass only, as already derived in [19]. In the neutralino sector, one finds that

$$\frac{1}{M_{\chi_1^\pm} M_{\chi_2^\pm}} \left\{ \prod_{i=1}^4 M_{\chi_i^0} - \tan^2 \theta_W \sin 2\beta M_W^2 \left(M_{\chi_1^\pm} M_{\chi_2^\pm} + \sin 2\beta M_W^2 \right) \right\} \times \left(\frac{g_2^9 g_3^{256/3}}{h_t^{27} h_b^{21} h_\tau^{10} g_1^{4099/33}} \right)^{1/61} \quad (3.15)$$

is an RG invariant. These invariants can be useful for determining the scale dependence of certain parameters from a few measured ones, and also they manifestly depict the correlation between the neutralino/chargino and gauge/Yukawa sectors. Finally, one notes that under the rescalings

$$\tilde{H}_{u,d} \rightarrow \left(\frac{g_2^9 g_3^{256/3}}{h_t^{27} h_b^{21} h_\tau^{10} g_1^{73/33}} \right)^{-1/122} \tilde{H}_{u,d}, \quad \tilde{B} \rightarrow g_1 \tilde{B}, \quad \tilde{W} \rightarrow g_2 \tilde{W} \quad (3.16)$$

the neutralino and chargino mass matrices become completely scale-invariant except for $SU(2)_L \otimes U(1)_Y$ breaking terms which mix Higgsinos and gauginos. This property could be useful in calculating and interpreting certain observables.

We have derived a number of RG invariants in the MSSM, and performed certain case studies for highlighting their phenomenological relevance. These invariants, as also emphasized in Introduction, could be quite useful for determining the origin of supersymmetry breaking, for testing the internal consistency of the model, and for obtaining certain sum rules that enable the prediction of certain unknown parameters from the known ones.

4. The RG invariants in U(1)' Model

The U(1)' extension of the MSSM is based on $SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)'$ gauge group with respective gauge couplings g_3, g_2, g_1 and g_1' . The Higgs sector is spanned by $\widehat{H}_u, \widehat{H}_d$ and \widehat{S} so that

$$\widehat{W}_{\text{Higgs}} = h_s \widehat{S} \widehat{H}_u \widehat{H}_d \tag{4.1}$$

is the unique superpotential comprising the Higgs superfields since U(1)' invariance (i) forbids the appearance of a bare μ parameter (as in the MSSM superpotential (3.1)), and (ii) does not allow for additional terms such as \widehat{S}^3 (as in the NMSSM superpotential to be discussed in the next section) [11, 12]. The full superpotential is obtained by adding (4.1) to the fermionic part given in (2.2).

The soft-breaking terms pertaining to Higgs and gaugino sectors are given by

$$\begin{aligned} -\mathcal{L}_{\text{soft}}^{\text{Higgs}} &= m_{H_u}^2 H_u^\dagger H_u + m_{H_d}^2 H_d^\dagger H_d + m_S^2 S^\dagger S + [h_s A_s S H_u \cdot H_d + \text{h.c.}], \\ -\mathcal{L}_{\text{soft}}^{\text{gaugino}} &= \frac{1}{2} \sum_{a=3,2,1,1'} [M_a \lambda_a \lambda_a + \text{h.c.}]. \end{aligned} \tag{4.2}$$

A comparison of the superpotential and soft-breaking terms with their MSSM counterparts (3.1) and (3.2) shows clearly the way the MSSM limit is reached. Indeed, below the U(1)' breaking scale the effective theory resembles the MSSM (it just resembles because, for instance, the neutralino sector of the MSSM is extended by U(1)' gaugino and singlino \widetilde{S} states) with the parameters

$$\mu_{\text{eff}} \equiv h_s \langle S \rangle, \quad \mu_{\text{eff}} B_{\text{eff}} \equiv h_s A_s \langle S \rangle \tag{4.3}$$

which are both stabilized at the weak scale as desired if the singlet develops a VEV $\langle S \rangle$ at the same scale [11, 12]. In essence, as far as the Higgs sector is concerned, the naturalness problem associated with the μ parameter of the MSSM is avoided as it is now generated dynamically by U(1)' breakdown [10].

The RGEs for model parameters are all listed in appendix A. For each quantity the way to MSSM limit is also described. Similar to the MSSM in section 3, one can construct a number of RG invariants by using the RGEs in appendix A. The invariants are tabulated in table 4. The first invariant I_1' is nothing but a direct generalization of the MSSM invariant I_1 . It expresses the fact that a specific combination of the inverse gauge coupling-squareds (with arbitrary c_1, c_2 and c_3) is independent of the energy scale.

The U(1)' model does not possess an exact RG invariant analogous to I_2 in the MSSM. The reason is that all four Yukawa couplings evolve with scale with their own RG equations; it is not possible form a scale-invariant combination of the Yukawa-squareds in the absence

Number	RG Invariant
I'_1	$\frac{3c_3 - c_2 - (33/5)c_1}{(6 + \rho\lambda_H)g_1^2} + \frac{c_1}{g_1^2} + \frac{c_2}{g_2^2} + \frac{c_3}{g_3^2}$
I'_2	$\approx h_s h_t^{3/7} h_b^{-39/7} h_\tau^{8/7} g_3^{64/7} g_2^{-9} g_1^{5/77} g_1'^{-a'}$
I'_3	$\frac{M_a}{g_a^2} \quad (a = 1', 1, 2, 3)$
I'_4	$\approx A_s + \frac{3}{7}A_t - \frac{39}{7}A_b + \frac{8}{7}A_\tau + \frac{64}{7}M_3 - 9M_2 + \frac{5}{77}M_1 - a'M'_1$
I'_5	$I_5 + \frac{\rho}{6 + \rho\lambda_H}(Q_E^2 - 2Q_L^2) M'_1 ^2$
I'_6	$I_6 - \frac{1}{2}m_S^2 + \frac{\rho}{6 + \rho\lambda_H}(Q_{H_u}^2 - \frac{1}{2}Q_S^2 - \frac{3}{2}Q_U^2) M'_1 ^2$
I'_7	$I_7 - \frac{1}{2}m_S^2 + \frac{\rho}{6 + \rho\lambda_H}(Q_{H_d}^2 - \frac{1}{2}Q_S^2 - \frac{3}{2}Q_D^2 - Q_L^2) M'_1 ^2$
I'_8	$I_8 + \frac{\rho}{6 + \rho\lambda_H}(Q_U^2 + Q_D^2 - 2Q_Q^2) M'_1 ^2$
I'_9	$I_9 + \frac{\rho}{6 + \rho\lambda_H}Q_Q^2 M'_1 ^2$
I'_{10}	$I_{10} + \frac{\rho}{6 + \rho\lambda_H}Q_U^2 M'_1 ^2$
I'_{11}	$I_{11} + \frac{\rho}{6 + \rho\lambda_H}Q_D^2 M'_1 ^2$
I'_{12}	$I_{12} + \frac{\rho}{6 + \rho\lambda_H}Q_L^2 M'_1 ^2$
I'_{13}	$I_{13} + \frac{\rho}{6 + \rho\lambda_H}Q_E^2 M'_1 ^2$
I'_{14}	$I_{14} - m_S^2 + \frac{\rho}{6 + \rho\lambda_H}(Q_{H_u}^2 + Q_{H_d}^2 - Q_S^2 - 3Q_Q^2 - Q_L^2) M'_1 ^2$
I'_{15}	$I_{15} - \frac{1}{2}m_S^2 + \frac{\rho}{6 + \rho\lambda_H}(Q_{H_d}^2 - \frac{1}{2}Q_S^2 - \frac{3}{2}Q_D^2 - \frac{3}{2}Q_L^2 + \frac{1}{4}Q_E^2) M'_1 ^2$
I'_{16}	$I_{16} + \frac{\rho}{6 + \rho\lambda_H}(2Q_Q^2 + Q_U^2 + Q_D^2) M'_1 ^2$
I'_{17}	$I_{17} + \frac{\rho}{6 + \rho\lambda_H}(Q_L^2 + \frac{1}{2}Q_E^2) M'_1 ^2$

Table 4: The RG invariant combinations of rigid and soft parameters in $U(1)'$ models (c_1, \dots, c_3 are arbitrary constants). Here I'_2 and I'_4 are approximate invariants derived in the text. The invariants constructed from scalar mass-squareds are written in terms of the MSSM invariants in table 1. The modifications are twofold: First, the MSSM invariants involving Higgs mass-squareds are shifted by the singlet mass-squared. Next, each invariant receives new contributions proportional to $|M'_1|^2$. Note that invariants pertaining to the first and second generations generically involve a single sfermion mass-squared since trilinear couplings do not contribute to their RGEs. For obtaining RG invariants containing a specific set of parameters it is necessary to form appropriate combinations of these tabulated ones, as exemplified in the text by a couple of case studies.

of a fifth equation that involves the same couplings (as $d \ln \mu / dt$ does). However, it is still possible to extract some important information about the UV/IR behaviors of the Yukawa couplings from their RGEs. Indeed, one can show that

$$\begin{aligned}
 \frac{h_s(Q_2)}{h_s(Q_1)} &= \left(\frac{h_t(Q_1)}{h_t(Q_2)} \right)^{3/7} \left(\frac{h_b(Q_1)}{h_b(Q_2)} \right)^{39/7} \left(\frac{h_\tau(Q_1)}{h_\tau(Q_2)} \right)^{8/7} \times \\
 &\times \left(\frac{g_3(Q_1)}{g_3(Q_2)} \right)^{64/7} \left(\frac{g_2(Q_1)}{g_2(Q_2)} \right)^9 \left(\frac{g_1(Q_1)}{g_1(Q_2)} \right)^{5/77} \left(\frac{g'_1(Q_2)}{g'_1(Q_1)} \right)^{a'} \times \\
 &\times \exp \left[-\frac{186}{7} \int_{t_{Q_1}}^{t_{Q_2}} dt' h_b^2(t') \right]
 \end{aligned} \tag{4.4}$$

with a' being a function of the $U(1)'$ charges

$$a' = \frac{\rho}{42 + 7\rho\lambda_H} (39Q_D^2 - 8Q_E^2 + 24Q_{H_d}^2 - 10Q_{H_u}^2 - 8Q_L^2 + 36Q_Q^2 - 7Q_S^2 - 3Q_U^2), \tag{4.5}$$

where ρ and λ_H are defined in appendix A. The importance of (4.4) stems from the fact that it explicitly expresses h_s in terms of the gauge couplings and rest of the Yukawas. This is important for both model building and phenomenological purposes since, in general, one has no information about the IR and UV values of h_s (in contrast to $h_{t,b,\tau}$ whose values at $Q = M_Z$ are known up to the ratio of the doublet VEVs). The ratio of the IR value of h_s to its UV value depends on all values of $h_b(t)$ in between because of the integration over h_b^2 at the right-hand side. However, this dependence is an extremely weak effect in low $\tan\beta$ domain where $-(186/7) \int_{t_{Q_1}}^{t_{Q_2}} dt' h_b^2(t') \sim 10^{-3}$. Presently, experiments have not shown yet a preferred interval for $\tan\beta$: it can range from 1 to m_t/m_b . However, in $U(1)'$ models (and also NMSSM to be discussed in the next section) large (and thus fine-tuned) $\tan\beta$ regime is not particularly preferred or needed to agree with the LEP bounds [29]. In fact, as demonstrated in [30], the $U(1)'$ models with a secluded sector naturally realize $\tan\beta \sim \mathcal{O}(1)$ with a heavy enough Z' gauge boson. Consequently, the lesson to be drawn from (4.4) is that given IR and UV values of the gauge and Yukawa couplings then the ratio $h_s(\text{IR})/h_s(\text{UV})$ is completely determined to an excellent approximation (the validity of which depends on how small $\tan\beta$ is). This conclusion enables us to introduce an approximate RG invariant

$$I'_2 \approx h_s h_t^{3/7} h_b^{-39/7} h_\tau^{8/7} g_3^{64/7} g_2^{-9} g_1^{5/77} g_1^{-a'} \quad (4.6)$$

which exhibits a rather weak scale-dependence especially when the VEVs of the two Higgs doublets are split within an $\mathcal{O}(1)$ factor. The invariant I'_2 given in table 4 is thus an approximate (albeit almost exact in low $\tan\beta$ domain) RG invariant.

The $U(1)'$ models possess an invariant like I_3 in the MSSM (see I'_3 in table 4). Indeed, ratio of a gaugino mass to the same gauge group's fine structure constant is an exact RG invariant. As in the MSSM, such invariants enable one to determine and predict gaugino mass of a given gauge group when others are given.

The RGEs of the trilinear couplings, given in appendix A, do not form an exact RG invariant for the reasons valid for Yukawa couplings. However, one can still establish correlations among the trilinears in order to extract information about their UV and IR behaviors. For example, the difference between the UV and IR values of A_s is related to those of the other parameters via

$$\begin{aligned} A_s(Q_2) - A_s(Q_1) = & -\frac{372}{7} \int_{t_{Q_1}}^{t_{Q_2}} dt' h_b^2(t') A_b(t') + \frac{3}{7} (A_t(Q_1) - A_t(Q_2)) + \\ & + \frac{39}{7} (A_b(Q_2) - A_b(Q_1)) + \frac{8}{7} (A_\tau(Q_1) - A_\tau(Q_2)) + \\ & + \frac{64}{7} M_3(Q_1) \left(\frac{g_3(Q_2)^2}{g_3(Q_1)^2} - 1 \right) - 9 M_2(Q_1) \left(\frac{g_2(Q_2)^2}{g_2(Q_1)^2} - 1 \right) + \\ & + \frac{5}{77} M_1(Q_1) \left(\frac{g_1(Q_2)^2}{g_1(Q_1)^2} - 1 \right) - a' M'_1(Q_1) \left(\frac{g'_1(Q_2)^2}{g'_1(Q_1)^2} - 1 \right) \quad (4.7) \end{aligned}$$

which depends on all values of $h_b^2 A_b$ in between the UV and IR scales. This dependence, however, is quite weak in low $\tan\beta$ domain (which is quite natural and does not pose

any difficulty with experimental bounds for the model under concern), and one can safely neglect this contribution. Then, to a good approximation (validity of which depends on how small $\tan\beta$ is) one can form an RG invariant

$$I'_4 \approx A_s + \frac{3}{7}A_t - \frac{39}{7}A_b + \frac{8}{7}A_\tau + \frac{64}{7}M_3 - 9M_2 + \frac{5}{77}M_1 - a'M'_1 \quad (4.8)$$

as is listed in table 4.

The RGEs for scalar soft mass-squareds depend on $S = \text{Tr}[Ym^2]$ and $S' = \text{Tr}[Qm^2]$ whose explicit expression is given in appendix A. In case soft masses are universal at some scale Q then S vanishes at all scales due to the absence of hypercharge-graviton-graviton anomaly. In this sense, as was discussed in detail in Sec. 3 when analyzing the MSSM RGEs, S is a viable probe of universality paradigm. These properties, however, cannot be continued to S' since even if the soft masses are universal S' does not need to vanish because $U(1)'$ charges are not guaranteed to cancel the gravitational anomaly of Z' boson. Indeed, the $U(1)'$ model is generically anomalous in that even if gravitational anomaly of $U(1)'$ is cancelled there remain all sorts of anomalies ($U(1)'^3$, $U(1)'SU(2)^2$, $U(1)'SU(3)_C^2$, ...) to be cancelled. These anomalies cannot be cancelled unless one introduces some exotic matter multiplets which necessarily disrupt the unification of gauge couplings [12, 31]. (It is worthy of noting that the model proposed in [32] extends the MSSM with a number of singlet chiral superfields, and determines the singlet $U(1)'$ charges by imposing anomaly cancellation.) Another option, as has recently been pointed out, is to introduce family non-universal $U(1)'$ charges for cancelling anomalies with minimal matter content [33]. Both options are beyond the scope of this work which explores RG invariant observables in minimal $U(1)'$ extension of the MSSM. Besides this, scale dependence of S' involves all soft masses, gaugino masses as well as trilinear couplings; it is not as compact as (3.7). This continues to be true unless $U(1)'$ charges of opposite-chirality same-flavor fermions obey the same ratios as the hypercharge. Moreover, $U(1)'$ charges of Higgs fields should exhibit a specific proportionality with their hypercharges. In what follows, we leave aside the question of anomalies and specific representations for $U(1)'$ charges, and simply take $S' \equiv 0$ at all scales of interest. (Within specific $U(1)'$ models such as the ones coming from $E(6)$ breaking or family non-universal $U(1)'$ models the probing power of S' can be analyzed explicitly.) With this simplifying assumption the RG invariant combinations of the soft masses $I'_5 - I'_{17}$, in parallel and with respect to those in the MSSM, are listed in table 4. The modifications in the MSSM invariants are twofold: First of all, each invariant picks up an additional contribution proportional to $|M'_1|^2$ (there would be an additional term from dS'/dt if S' were not taken vanishing). The proportionality constant involves $U(1)'$ beta function and a linear combination of charge-squareds with coefficients identical to those of the soft mass-squareds relevant for the invariant under consideration. The other modification in MSSM invariants concerns the presence of Higgs mass-squareds. Indeed, if an MSSM invariant involves $m_{H_u}^2$ or $m_{H_d}^2$ then the corresponding $U(1)'$ invariant is necessarily shifted by $-(1/2)m_S^2$. The reason for this is the presence of terms proportional to h_s^2 in the beta functions of $m_{H_u}^2$, $m_{H_d}^2$ and m_S^2 .

A couple of case studies can shed light on certain aspects of the RG invariants in $U(1)'$ models. First, let us consider the RG invariant I'_8 in table 4. Similar to its MSSM analogue (3.10), it gives rise to

$$\begin{aligned}
 m_{\tilde{t}_R}^2(Q_2) + m_{\tilde{b}_R}^2(Q_2) - 2m_{\tilde{t}_L}^2(Q_2) &= 3|M_2(Q_1)|^2 \left[\left(\frac{g_2(Q_2)}{g_2(Q_1)} \right)^4 - 1 \right] - \\
 &\quad - \frac{1}{11} |M_1(Q_1)|^2 \left[\left(\frac{g_1(Q_2)}{g_1(Q_1)} \right)^4 - 1 \right] + \\
 &\quad + \frac{\rho}{6 + \rho\lambda_H} (Q_U^2 + Q_D^2 - 2Q_Q^2) \times \\
 &\quad \times |M'_1(Q_1)|^2 \left[\left(\frac{g'_1(Q_2)}{g'_1(Q_1)} \right)^4 - 1 \right] \quad (4.9)
 \end{aligned}$$

when soft mass-squareds are all universal at some scale $Q = Q_1$. This relation is independent of the Higgs sector parameters; it is sensitive to only the isospin and Abelian group factors. In fact, it feels whether the gauge sector is minimal or not by the inclusion of the corresponding gaugino mass in the sum rule. Therefore, via the last term $\propto |M'_1|^2$, it obtains the potential of probing the existence of an additional $U(1)'$ gauge invariance provided that one can perform precise measurements and consistency checks with other sectors of the theory. Of course, (4.9) can be used to establish a relation between the stop and sbottom mixing angles in the same spirit as (3.11).

One notices that it is not possible to construct an RG invariant which feels only the extensions in the Higgs sector. The reason is that in a given sum rule each mass-squared parameter is accompanied by an additional term $\propto Q^2 |M'_1|^2$ any attempt at cancelling terms involving $|M'_1|^2$ necessarily ends up with cancelling m_S^2 contribution. Of course, within a specific representation for $U(1)'$ invariance charges of various fields could be correlated to cancel out without nullifying the coefficient of m_S^2 in the final sum rule. It was this property of hypercharge symmetry that allowed us to arrive at (3.12) in the MSSM section above. For instance, if Q_E^2 happens to be proportional to $Q_{H_u}^2 + Q_{H_d}^2 - Q_S^2 - 3Q_Q^2 - Q_L^2$ then I'_{14} and I'_{13} can be used to relate mass-squareds of Higgs fields to those of sfermions and MSSM gauginos.

There exist certain RG-invariant combinations of the soft-mass squareds which depend on extensions in neither the gauge nor the Higgs sectors. Invariants of this kind can be easily constructed by linearly combining those in table 4. For instance,

$$I'_8 + 2I'_9 - I'_{10} - I'_{11} = \left(m_{\tilde{t}_R}^2 - m_{\tilde{u}_R}^2 \right) + \left(m_{\tilde{b}_R}^2 - m_{\tilde{d}_R}^2 \right) - 2 \left(m_{\tilde{t}_L}^2 - m_{\tilde{u}_L}^2 \right) \quad (4.10)$$

is an RG invariant in both MSSM and $U(1)'$ models. Clearly, this kind of quantities are completely insensitive to modifications in the Higgs and gauge sectors; they exclusively probe the sfermion sector.

In general, within specific string or supergravity models, the soft parameters exhibit various interrelations which give rise to a spectrum of discriminative values for the RG invariants. This can be used for predicting what specific model could be responsible for

supersymmetry breaking. Indeed, as one recalls from discussions of the MSSM invariants within no-scale supergravity, dilaton-domination and flux-induced soft terms, measurements of RG invariants could be a useful tool for determining which high scale model is operating as more and more data accumulate. For the $U(1)'$ model under concern it is convenient to discuss the predictive powers of RG invariants with respect to the MSSM ones tabulated in tables 2 and 3: first of all, if the UV scale model realizes a universal gaugino mass M then each of $I'_5/M^2 - I'_{17}/M^2$ is shifted by an amount

$$\frac{\rho}{6 + \rho\lambda_H} (a_X Q_X^2 + b_Y Q_Y^2 + c_Z Q_Z^2 + \dots) \quad (4.11)$$

if the invariant under concern is composed of $a_X m_X^2 + b_Y m_Y^2 + c_Z m_Z^2 + \dots$. If an invariant does not contain Higgs soft mass-squareds or $m_0 = 0$ for the model under concern then this is the only modification in an RG invariant with respect to its MSSM value given in tables 2 and 3. On the other hand, if an invariant consists of the Higgs masses, e.g. I'_6, I'_7, I'_{14} , then departure from the MSSM expression occurs in both m_0^2 and M^2 directions. The discriminative power of an RG invariant depends on its correlation with others for a given GUT-scale configuration. Generically, if the mass-squareds of Higgs doublets are present in an invariant so is that of the Higgs singlet.

The RG invariants do have interesting implications also for chargino and neutralino sectors of the model. For instance, from the product of two chargino masses one immediately finds that the quantity

$$\frac{1}{\langle S \rangle} \left(M_{\chi_1^\pm} M_{\chi_2^\pm} + M_W^2 \sin 2\beta \right) \left(h_t^{3/7} h_b^{-39/7} h_\tau^{8/7} g_3^{64/7} g_2^{-11} g_1^{5/77} g_1^{-a'} \right) \quad (4.12)$$

is an approximate RG invariant. This RG invariant differs from its MSSM analogue (3.14) by modifications in powers of the gauge and Yukawa couplings and by the presence of the singlet VEV $\langle S \rangle$. The presence of the singlet VEV stems from the fact that the μ parameter in the MSSM is generated dynamically by the singlet VEV: $\mu_{\text{eff}} = h_s \langle S \rangle$.

The neutralino sector is sensitive to both the Higgs singlet and $U(1)'$ gaugino. First of all, sum of the neutralino masses obey

$$\sum_{i=1}^6 M_{\chi_i^0} = M'_1 + M_1 + M_2 = \left(g_1'^2 + g_1^2 + g_2^2 \right) \frac{M_{1/2}}{g_0^2} \quad (4.13)$$

when the gaugino masses unify into $M_{1/2}$ at the scale where gauge couplings do into g_0 . The sum of the squared-masses of neutralinos depend on both M'_1 and $h_s \langle S \rangle$. Therefore, a correlated analysis of neutralino and chargino sectors provide important information on whether the MSSM is extended by new gauge symmetries and/or new Higgs representations. The neutralino sector admits several sum/product rules similar to (3.15) in the MSSM, and they can be used to form novel RG invariant combinations of the chargino/neutralino parameters in the same spirit as (4.13) and (4.12). One keeps in mind, however, that invariants involving the Higgs singlet is always approximate in the sense of (4.4).

In this section we have derived a number of RG invariants in $U(1)'$ models, and performed certain case studies for highlighting their phenomenological relevance. These invariants (albeit approximate for Yukawa couplings and trilinear soft terms) could be useful for establishing gauge and/or Higgs extension with respect to the MSSM.

5. The RG invariants in the NMSSM

The next-to-minimal supersymmetric model possesses no gauge extension with respect to MSSM. Its Higgs sector is spanned by \widehat{H}_u , \widehat{H}_d and \widehat{S} so that

$$\widehat{W}_{\text{Higgs}} = h_s \widehat{S} \widehat{H}_u \widehat{H}_d + \frac{k_s}{6} \widehat{S}^3 \quad (5.1)$$

and

$$\begin{aligned} -\mathcal{L}_{\text{soft}}^{\text{Higgs}} &= m_{H_u}^2 H_u^\dagger H_u + m_{H_d}^2 H_d^\dagger H_d + m_S^2 S^\dagger S + \left[h_s A_s S H_u \cdot H_d + \frac{k_s}{6} A_k S^3 + \text{h.c.} \right], \\ -\mathcal{L}_{\text{soft}}^{\text{gaugino}} &= \frac{1}{2} \sum_{a=3,2,1} [M_a \lambda_a \lambda_a + \text{h.c.}], \end{aligned} \quad (5.2)$$

where the singlet cubic interaction in the superpotential is needed to generate a potential for S (this field does not have a D-term support to obtain a potential). The induction of effective μ and B parameters are similar to those of the $U(1)'$ model given in (4.3). The main difference from the $U(1)'$ model lies in the fact that the \widehat{S} is a pure singlet (in both MSSM and NMSSM) so that it is allowed to develop a cubic interaction in the superpotential.

The RGEs of the rigid and soft parameters of the model are all listed in appendix B. We also discuss the MSSM limits of individual RGEs for easy comparison of the corresponding RG invariants. The RG-invariant quantities in the model are listed in table 5. Obviously, the RG invariant combinations of gauge couplings, I_1'' , remain the same as in the MSSM.

In close similarity to $U(1)'$ models, the Yukawa couplings do not possess an exact RG invariant. However, it is still possible to express one of the Yukawas in terms of the rest and gauge couplings. For instance, the singlet cubic coupling is related to others via

$$\begin{aligned} \frac{k_s(Q_2)}{k_s(Q_1)} &= \left(\frac{h_t(Q_1)}{h_t(Q_2)} \right)^{3/7} \left(\frac{h_b(Q_1)}{h_b(Q_2)} \right)^{15/7} \left(\frac{h_\tau(Q_2)}{h_\tau(Q_1)} \right)^{6/7} \left(\frac{h_s(Q_2)}{h_s(Q_1)} \right)^3 \times \\ &\times \left(\frac{g_3(Q_2)}{g_3(Q_1)} \right)^{832/189} \left(\frac{g_2(Q_2)}{g_2(Q_1)} \right)^{27/7} \left(\frac{g_1(Q_2)}{g_1(Q_1)} \right)^{23/77} \times \\ &\times \exp \left[\frac{197}{7} \int_{t_{Q_1}}^{t_{Q_2}} dt' h_b^2(t') \right]. \end{aligned} \quad (5.3)$$

The importance of this relation stems from the fact that it explicitly expresses k_s in terms of the gauge couplings and rest of the Yukawas. This is important for both model building and phenomenological purposes since, in general, one has no information about the IR and UV values of both h_s and k_s (in contrast to $h_{t,b,\tau}$ whose values at $Q = M_Z$ are known up to the ratio of the doublet VEVs), and it is advantageous to know at least one's value in terms of the rest. The ratio of the IR value of k_s to its UV value depends on all values of $h_b(t)$ in between because of the integration over h_b^2 at the right-hand side. However, this dependence is an extremely weak effect in low $\tan \beta$ domain where $(197/7) \int_{t_{Q_1}}^{t_{Q_2}} dt' h_b^2(t') \sim 10^{-3}$. Presently, experiments have not shown yet a preferred interval for $\tan \beta$: it can range from

Number	RG Invariant
I''_1	$\frac{c_1}{g_1^2} + \frac{c_2}{g_2^2} + \frac{33c_1+5c_2}{15g_3^2}$
I''_2	$\approx k_s h_t^{3/7} h_b^{15/7} h_\tau^{-6/7} h_s^{-3} g_3^{-832/189} g_2^{-27/7} g_1^{-23/77}$
I''_3	$\frac{M_a}{g_a^2} (a = 1, 2, 3)$
I''_4	$\approx A_k + \frac{3}{7}A_t + \frac{15}{7}A_b - \frac{6}{7}A_\tau - \frac{382}{189}M_3 - \frac{27}{7}M_2 - \frac{23}{77}M_1$
I''_5	I_5
I''_8	I_8
I'_9	I_9
I''_{10}	I_{10}
I''_{11}	I_{11}
I''_{12}	I_{12}
I''_{13}	I_{13}
I''_{16}	I_{16}
I''_{17}	I_{17}

Table 5: The RG invariant combinations of rigid and soft parameters in the NMSSM (c_1 and c_2 are arbitrary constants). Here I'_2 and I'_4 are approximate invariants derived in the text. The invariants constructed from scalar mass-squareds are written in terms of the MSSM invariants in table 1. The missing rows (with respect to table 1) indicate that there are no analogous RG invariant combinations of the scalar soft mass-squareds (the ones that depend on the Higgs sector parameters).

1 to m_t/m_b . However, in NMSSM large (and thus fine-tuned) $\tan \beta$ regime is not particularly preferred or needed to explain the LEP limits [34]. Consequently, (5.3) implies that, given IR and UV values of the gauge and Yukawa couplings, then the ratio $k_s(\text{IR})/k_s(\text{UV})$ is completely determined to an excellent approximation (the validity of which depends on how small $\tan \beta$ is). This conclusion enables us to introduce an approximate RG invariant

$$I''_2 \approx k_s h_t^{3/7} h_b^{15/7} h_\tau^{-6/7} h_s^{-3} g_3^{-832/189} g_2^{-27/7} g_1^{-23/77} \tag{5.4}$$

which exhibits a rather weak scale-dependence especially when the VEVs of the two Higgs doublets are split within an $\mathcal{O}(1)$ factor [34]. The invariant I''_2 given in table 5 is thus an approximate RG invariant.

The ratio of the gaugino masses to the corresponding fine structure constant, I''_3 in table 5, is an RG invariant, and it equals the corresponding invariant in the MSSM.

The behaviors of the trilinear couplings are similar to Yukawas. They do not admit an exact RG invariant. However, one can correlate their UV and IR values as in the U(1)' models. For example, the difference between the UV and IR values of A_k is related to those of the other parameters via

$$A_k(Q_2) - A_k(Q_1) = \frac{384}{7} \int_{t_{Q_1}}^{t_{Q_2}} dt' h_b^2(t') A_b(t') + \frac{3}{7} (A_t(Q_1) - A_t(Q_2)) + \frac{15}{7} (A_b(Q_1) - A_b(Q_2)) +$$

$$\begin{aligned}
 & + \frac{6}{7} (A_\tau(Q_2) - A_\tau(Q_1)) + 3 (A_s(Q_2) - A_s(Q_1)) - \\
 & - \frac{382}{189} M_3(Q_1) \left(\frac{g_3(Q_2)^2}{g_3(Q_1)^2} - 1 \right) - \frac{27}{7} M_2(Q_1) \left(\frac{g_2(Q_2)^2}{g_2(Q_1)^2} - 1 \right) - \\
 & - \frac{23}{77} M_1(Q_1) \left(\frac{g_1(Q_2)^2}{g_1(Q_1)^2} - 1 \right) \tag{5.5}
 \end{aligned}$$

which depends on all values of $h_b^2 A_b$ in between the UV and IR scales. This dependence, however, is quite weak in low $\tan \beta$ domain, and as experiments are not pushing for high $\tan \beta$ regime for the NMSSM, this dependence on $h_b^2 A_b$ can safely be neglected. Then, in low $\tan \beta$ regime, one can form an approximate RG invariant

$$I'_4 \approx A_k + \frac{3}{7} A_t + \frac{15}{7} A_b - \frac{6}{7} A_\tau - \frac{382}{189} M_3 - \frac{27}{7} M_2 - \frac{23}{77} M_1 \tag{5.6}$$

as listed in table 5.

As in the MSSM and $U(1)'$ models the soft squared-mass parameters do also form a number of RG invariants. These are listed in table 5. Perhaps, the most interesting aspect of the NMSSM is that its Higgs sector parameters do not admit any RG invariant. The reason is that the RG running of m_S^2 is necessarily affected by the cubic singlet coupling via $(3m_S^2 + |A_k|^2)k_s^2$ whereas running of the squared-masses of other fields do not involve terms $\propto k_s^2$. Hence, this term cannot be cancelled to form an invariant, and therefore, the MSSM RG invariants I_6 , I_7 , I_{14} , and I_{15} (which consist of the Higgs squared-masses) in table 1 do not possess any analogue in table 5. Physically, this is related to the fact that neither F terms nor soft terms generate operators of the form $k_s^2 |S|^2 (|H_u|^2, |H_d|^2, |\tilde{Q}|^2, \dots)$. It is convenient to dwell on this point by examining one of the would-be invariants. For instance, in the present model I_{14} in table 1 generalizes to

$$\frac{d}{dt} (I_{14} - m_S^2) = - \left(3m_S^2 + |A_k|^2 \right) k_s^2 \tag{5.7}$$

so that $I_{14} - m_S^2$ is not a scale-invariant observable; it exhibits a nontrivial RG flow unless (i) $k_s = 0$ or (ii) $m_S^2 = -|A_k|^2/3$. The former is disfavored for it gives rise to a flat direction for S [13]. The latter, however, represents a fixed point solution for m_S^2 in that at the scale it holds m_S^2 is guaranteed to be negative and hence the theory below $|A_k|$ generates the MSSM as an effective theory. Clearly, if $|A_k| \sim \mathcal{O}(\text{TeV})$ the MSSM Higgs sector gets correctly stabilized at the desired scale.

Looking from a different angle, (5.7) provides an experimental testing ground (presumably after $LHC \oplus ILC$) for knowing if the model under concern is NMSSM or $U(1)'$ extension of the MSSM. Indeed, in $U(1)'$ models the right hand side of (5.7) is $\propto g_1'^2 |M_1'|^2$ and it can be written as a total derivative to form the invariant I'_{14} in table 4. Moreover, as depicted in table 4 all RG invariants of soft mass-squareds systematically contain $|M_1'|^2$ so that after sufficient number of precise measurements one can make sure if the model under concern involves a new gaugino or not. In contrast to this, the right hand side of (5.7) cannot be written as a total derivative; moreover, it shows up only in those would-be in-

variants which include the Higgs soft-mass squareds. The rest of the invariants, as shown in table 5, are identical to those in the MSSM. In this sense, non-invariance of the Higgs sector parameters can provide a viable signal of NMSSM in future collider tests.

The RG-invariant combinations of the squared soft masses listed in table 5 give rise to certain correlations or sum rules which are identical to those derived in the MSSM. For instance, I''_8 relates stop plus sbottom splittings to the isospin and hypercharge gaugino masses in the same way as (3.10).

The RG invariant combinations of the chargino/neutralino systems are similar to ones in $U(1)'$ models. Indeed, (4.12) now becomes

$$\frac{1}{\langle S \rangle} \left(M_{\chi_1^\pm} M_{\chi_2^\pm} + M_W^2 \sin 2\beta \right) \left(h_t^{-1/7} h_b^{-5/7} h_\tau^{2/7} g_3^{832/567} g_2^{-5/7} g_1^{23/231} \right) \quad (5.8)$$

is an approximate RG invariant in the sense of (5.3). The singlet VEV $\langle S \rangle$ arises due to the dynamical origin of the MSSM μ parameter: $\mu_{\text{eff}} = h_s \langle S \rangle$.

The neutralino sector is interesting in that sum of the neutralino masses satisfy

$$\sum_{i=1}^5 M_{\chi_i^0} = M_1 + M_2 = (g_1^2 + g_2^2) \frac{M_{1/2}}{g_0^2} \quad (5.9)$$

which is identical to the MSSM prediction. The NMSSM effects show up when we consider sum of the neutralino mass-squareds or when we consider their products. Such quantities, too, can be expressed in terms of the RG invariants at low values of $\tan \beta$. Their validity and construction are not different than (5.8).

In this section we have analyzed the RGEs of the NMSSM for determining RG invariant combinations of the lagrangian parameters. Concerning the scale dependencies of the Yukawa couplings and trilinear soft terms, the behavior is similar to $U(1)'$ model. On the other hand, RG invariants made up of gauge couplings and scalar soft mass-squareds are the same as in the MSSM. The model radically differs from the MSSM and $U(1)'$ model due to the absence of RG invariants containing the Higgs mass-squareds.

6. Conclusion

In this work, using one loop RGEs, we have derived a number of scale-invariant observables in softly-broken supersymmetric models, and illustrated their phenomenological implications by various case studies. We have first studied the MSSM and then its minimal extensions, $U(1)'$ models and NMSSM, in a comparative manner.

In general, each supersymmetric model possesses RG invariants in gauge, Yukawa and soft-breaking sectors. The invariants of the MSSM, of $U(1)'$ model and of the NMSSM are listed in tables 1, 4 and 5, respectively. In general, RG invariants vary from model to model though those associated exclusively with their common part, the sfermion sector, may be combined to obtain invariants valid for all three models (see e.g. the combination 4.10).

The RG-invariant combinations of the gauge couplings and gaugino masses are identical for the MSSM and NMSSM whereas additional gauge coupling and mass of the associated gaugino introduces additional terms for the $U(1)'$ model. For phenomenological purposes, such invariants prove particularly useful when gauge couplings and gaugino masses unify at high scale.

The μ parameter, Yukawa couplings and gauge couplings combine to form an RG invariant in the MSSM. This, however, is not the case in $U(1)'$ models and NMSSM. In these models, the best one can do is to correlate one of the Yukawas in terms of the rest so that an approximate RG invariant emerges within a specific domain of the parameter space. In fact, I_2' in table 4 and I_2'' in table 5 serve as RG invariants only for low values of $\tan\beta$.

The Higgs bilinear soft mass B and sfermion-sfermion-Higgs trilinear couplings form an exact RG invariant in the MSSM. However, for $U(1)'$ models and NMSSM there are no such invariants, and as for the Yukawa sector, all one can do is to realize an approximate invariant in a specific domain of the parameter space. In fact, I_4' in table 4 and I_4'' in table 5 behave as RG invariants only at low values of $\tan\beta$.

The MSSM possesses a number of RG invariants containing the squared-masses of the scalars. They can be grouped into two classes: The ones that are not sensitive to whether the scalar masses attain a universal configuration and the ones that are sensitive (via the quantity S) to such a configuration. Therefore, the S dependence of the invariants serves as a tool for probing the UV scale correlations of the soft mass-squareds (as part of the minimal supergravity configuration). Moreover, as shown in tables 3 and 4, the invariants take on a specific set of values for each mechanism of supersymmetry breaking, and therefore, they can be used for determining the origin of supersymmetry breaking.

The RG invariants of scalar mass-squareds in the MSSM get modified by the $U(1)'$ gaugino mass and by the singlet mass-squared. In particular, those MSSM invariants which depend on the Higgs mass-squareds are generically shifted by the singlet mass-squared. It is possible to form new invariants that involve only the $U(1)'$ gaugino mass. On the other hand, invariants that depend only on the singlet mass-squared cannot be formed (unless one uses a specific representation for $U(1)'$ charges).

The situation in the NMSSM is interesting in that the Higgs mass-squareds cannot be combined to form an invariant because of the presence of cubic singlet coupling in the superpotential. This non-invariance itself can be useful for model identification at future collider studies. On the other hand, mass-squareds of scalar quarks and leptons admit RG-invariant configurations that are identical to those in the MSSM.

In general, the RG invariants are useful for both model-building and phenomenological purposes as they make various indirect relations manifest. This enhances one's knowledge of various dependencies and correlations among the model parameters. Moreover, they give rise to certain sum rules which can be quite useful for determining the underlying model and origin of supersymmetry breaking as data accumulate at detectors. Various relations require an accurate measurement of a subset of parameters which could be possible after $LHC\oplus ILC$.

Acknowledgments

The author thanks Lisa Everett for discussions on RGEs. This work was partially supported by Turkish Academy of Sciences through GEBIP grant, and by the Scientific and Technical Research Council of Turkey through project 104T503.

A. Renormalization group equations in U(1)' model

In this appendix we list down the RGEs for U(1)' models by extending [12] to cases with finite bottom and tau Yukawas in a way including all three generations of sfermions. The one-loop RGEs of the gauge couplings are given by

$$\begin{aligned}
 \frac{dg_3}{dt} &= (2N_F - 9) g_3^3 \\
 \frac{dg_2}{dt} &= (2N_F - 5) g_2^3 \\
 \frac{dg_1}{dt} &= \left(2N_F + \frac{3}{5}\right) g_1^3 \\
 \frac{dg'_1}{dt} &= (2N_F + \rho\lambda_H) g_1'^3,
 \end{aligned} \tag{A.1}$$

where $t \equiv (4\pi)^{-2} \ln Q/M_{\text{GUT}}$, $N_F = 3$, $\lambda_H = Q_{H_d}^2 + Q_{H_u}^2 + \frac{1}{2}Q_S^2$, and

$$\rho = \frac{4}{6Q_Q^2 + 3(Q_U^2 + Q_D^2) + 2Q_L^2 + Q_E^2} \tag{A.2}$$

which is obtained by requiring $g_a^2 \text{Tr}[Q^2]$ to be identical for all group factors. The U(1)' charges $Q_{H_u, \dots, E}$ are family-universal. The corresponding MSSM RGEs are recovered by setting $g'_1 = 0$.

The evolutions of the superpotential parameters are given by

$$\begin{aligned}
 \frac{dh_t}{dt} &= h_t \left(6h_t^2 + h_b^2 + h_s^2 - \frac{16}{3}g_3^2 - 3g_2^2 - \frac{13}{15}g_1^2 - \rho(Q_{H_u}^2 + Q_Q^2 + Q_U^2) g_1'^2\right) \\
 \frac{dh_b}{dt} &= h_b \left(6h_b^2 + h_t^2 + h_\tau^2 + h_s^2 - \frac{16}{3}g_3^2 - 3g_2^2 - \frac{7}{15}g_1^2 - \rho(Q_{H_d}^2 + Q_Q^2 + Q_D^2) g_1'^2\right) \\
 \frac{dh_\tau}{dt} &= h_\tau \left(4h_\tau^2 + 3h_b^2 + h_s^2 - 3g_2^2 - \frac{9}{5}g_1^2 - \rho(Q_{H_d}^2 + Q_L^2 + Q_E^2) g_1'^2\right) \\
 \frac{dh_s}{dt} &= h_s \left(4h_s^2 + 3h_t^2 + 3h_b^2 + h_\tau^2 - 3g_2^2 - \frac{3}{5}g_1^2 - \rho(Q_{H_d}^2 + Q_{H_u}^2 + Q_S^2) g_1'^2\right)
 \end{aligned} \tag{A.3}$$

which reduce to the corresponding RGEs in the MSSM after setting $g'_1 = 0$, identifying $d \ln h_s / dt$ with $d \ln \mu / dt$ in the last equation, and taking $h_s = 0$ everywhere else (since now a dynamical field \hat{S} does not exist at all).

The gaugino masses evolve as

$$\frac{dM_3}{dt} = (4N_F - 18) g_3^2 M_3$$

$$\begin{aligned}
 \frac{dM_2}{dt} &= (4N_F - 10) g_2^2 M_2 \\
 \frac{dM_1}{dt} &= \left(4N_F + \frac{6}{5}\right) g_1^2 M_1 \\
 \frac{dM'_1}{dt} &= (4N_F + \rho(2Q_{H_d}^2 + 2Q_{H_u}^2 + Q_S^2)) g_1'^2 M'_1
 \end{aligned} \tag{A.4}$$

which reduce to the RGEs in the MSSM after setting $M'_1 = 0$.

The RG evolutions of the trilinear couplings are given by

$$\begin{aligned}
 \frac{dA_t}{dt} &= 2(6h_t^2 A_t + h_b^2 A_b + h_s^2 A_s) + \\
 &\quad + 2\left(\frac{16}{3}g_3^2 M_3 + 3g_2^2 M_2 + \frac{13}{15}g_1^2 M_1 + \rho(Q_{H_u}^2 + Q_Q^2 + Q_U^2) g_1'^2 M'_1\right) \\
 \frac{dA_b}{dt} &= 2(6h_b^2 A_b + h_t^2 A_t + h_\tau^2 A_\tau + h_s^2 A_s) + \\
 &\quad + 2\left(\frac{16}{3}g_3^2 M_3 + 3g_2^2 M_2 + \frac{7}{15}g_1^2 M_1 + \rho(Q_{H_d}^2 + Q_Q^2 + Q_D^2) g_1'^2 M'_1\right) \\
 \frac{dA_\tau}{dt} &= 2(4h_\tau^2 A_\tau + 3h_b^2 A_b + h_s^2 A_s) + \\
 &\quad + 2\left(3g_2^2 M_2 + \frac{9}{5}g_1^2 M_1 + \rho(Q_{H_d}^2 + Q_L^2 + Q_E^2) g_1'^2 M'_1\right) \\
 \frac{dA_s}{dt} &= 2(4h_s^2 A_s + 3h_t^2 A_t + 3h_b^2 A_b + h_\tau^2 A_\tau) + \\
 &\quad + 2\left(3g_2^2 M_2 + \frac{3}{5}g_1^2 M_1 + \rho(Q_{H_d}^2 + Q_{H_u}^2 + Q_S^2) g_1'^2 M'_1\right),
 \end{aligned} \tag{A.5}$$

where RGEs of the corresponding quantities within the MSSM are obtained by $g'_1 = 0$, $h_s = 0$ and $A_s = B$, B being the Higgs soft bilinear coupling.

The scalar soft mass-squared parameters evolve according to

$$\begin{aligned}
 \frac{dm_{H_u}^2}{dt} &= 2\left(m_{H_u}^2 + m_{H_d}^2 + m_S^2 + |A_s|^2\right) h_s^2 + 6\left(m_{H_u}^2 + m_{t_L}^2 + m_{t_R}^2 + |A_t|^2\right) h_t^2 - \\
 &\quad - 8\left(\frac{3}{4}g_2^2 |M_2|^2 + \frac{3}{20}g_1^2 |M_1|^2 + \frac{1}{2}\rho Q_{H_u}^2 g_1'^2 |M'_1|^2\right) + \frac{3}{5}g_1^2 S + \rho Q_{H_u} g_1'^2 S' \\
 \frac{dm_{H_d}^2}{dt} &= 2\left(m_{H_u}^2 + m_{H_d}^2 + m_S^2 + |A_s|^2\right) h_s^2 + 2\left(m_{H_d}^2 + m_{\tau_L}^2 + m_{\tau_R}^2 + |A_\tau|^2\right) h_\tau^2 + \\
 &\quad + 6\left(m_{H_d}^2 + m_{t_L}^2 + m_{b_R}^2 + |A_b|^2\right) h_b^2 - \\
 &\quad - 8\left(\frac{3}{4}g_2^2 |M_2|^2 + \frac{3}{20}g_1^2 |M_1|^2 + \frac{1}{2}\rho Q_{H_d}^2 g_1'^2 |M'_1|^2\right) - \frac{3}{5}g_1^2 S + \rho Q_{H_d} g_1'^2 S' \\
 \frac{dm_S^2}{dt} &= 4\left(m_{H_u}^2 + m_{H_d}^2 + m_S^2 + |A_s|^2\right) h_s^2 - 4\rho Q_S^2 g_1'^2 |M'_1|^2 + \rho Q_S g_1'^2 S' \\
 \frac{dm_{t_L}^2}{dt} &= 2\left(m_{t_L}^2 + m_{H_d}^2 + m_{b_R}^2 + |A_b|^2\right) h_b^2 + 2\left(m_{t_L}^2 + m_{H_u}^2 + m_{t_R}^2 + |A_t|^2\right) h_t^2 - \\
 &\quad - 8\left(\frac{4}{3}g_3^2 |M_3|^2 + \frac{3}{4}g_2^2 |M_2|^2 + \frac{1}{60}g_1^2 |M_1|^2 + \frac{1}{2}\rho Q_Q^2 g_1'^2 |M'_1|^2\right) +
 \end{aligned}$$

$$\begin{aligned}
 & + \frac{1}{5}g_1^2 S + \rho Q_Q g_1'^2 S' \\
 \frac{dm_{\tilde{t}_R}^2}{dt} &= 4 \left(m_{\tilde{t}_L}^2 + m_{H_u}^2 + m_{\tilde{t}_R}^2 + |A_t|^2 \right) h_t^2 - \\
 & - 8 \left(\frac{4}{3}g_3^2 |M_3|^2 + \frac{4}{15}g_1^2 |M_1|^2 + \frac{1}{2}\rho Q_U^2 g_1'^2 |M_1'|^2 \right) - \frac{4}{5}g_1^2 S + \rho Q_U g_1'^2 S' \\
 \frac{dm_{\tilde{b}_R}^2}{dt} &= 4 \left(m_{\tilde{t}_L}^2 + m_{H_d}^2 + m_{\tilde{b}_R}^2 + |A_b|^2 \right) h_b^2 - \\
 & - 8 \left(\frac{4}{3}g_3^2 |M_3|^2 + \frac{1}{15}g_1^2 |M_1|^2 + \frac{1}{2}\rho Q_D^2 g_1'^2 |M_1'|^2 \right) + \frac{2}{5}g_1^2 S + \rho Q_D g_1'^2 S' \\
 \frac{dm_{\tilde{\tau}_L}^2}{dt} &= 2 \left(m_{\tilde{\tau}_L}^2 + m_{H_d}^2 + m_{\tilde{\tau}_R}^2 + |A_\tau|^2 \right) h_\tau^2 - \\
 & - 8 \left(\frac{3}{4}g_2^2 |M_2|^2 + \frac{3}{20}g_1^2 |M_1|^2 + \frac{1}{2}\rho Q_L^2 g_1'^2 |M_1'|^2 \right) - \frac{3}{5}g_1^2 S + \rho Q_L g_1'^2 S' \\
 \frac{dm_{\tilde{\tau}_R}^2}{dt} &= 4 \left(m_{\tilde{\tau}_L}^2 + m_{H_d}^2 + m_{\tilde{\tau}_R}^2 + |A_\tau|^2 \right) h_\tau^2 - \\
 & - 8 \left(\frac{3}{5}g_1^2 |M_1|^2 + \frac{1}{2}\rho Q_E^2 g_1'^2 |M_1'|^2 \right) + \frac{6}{5}g_1^2 S + \rho Q_E g_1'^2 S' \\
 \frac{dm_{\tilde{u}_L}^2}{dt} &= -8 \left(\frac{4}{3}g_3^2 |M_3|^2 + \frac{3}{4}g_2^2 |M_2|^2 + \frac{1}{60}g_1^2 |M_1|^2 + \frac{1}{2}\rho Q_Q^2 g_1'^2 |M_1'|^2 \right) + \\
 & + \frac{1}{5}g_1^2 S + \rho Q_Q g_1'^2 S' \\
 \frac{dm_{\tilde{u}_R}^2}{dt} &= -8 \left(\frac{4}{3}g_3^2 |M_3|^2 + \frac{4}{15}g_1^2 |M_1|^2 + \frac{1}{2}\rho Q_U^2 g_1'^2 |M_1'|^2 \right) - \frac{4}{5}g_1^2 S + \rho Q_U g_1'^2 S' \\
 \frac{dm_{\tilde{d}_R}^2}{dt} &= -8 \left(\frac{4}{3}g_3^2 |M_3|^2 + \frac{1}{15}g_1^2 |M_1|^2 + \frac{1}{2}\rho Q_D^2 g_1'^2 |M_1'|^2 \right) + \frac{2}{5}g_1^2 S + \rho Q_D g_1'^2 S' \\
 \frac{dm_{\tilde{e}_L}^2}{dt} &= -8 \left(\frac{3}{4}g_2^2 |M_2|^2 + \frac{3}{20}g_1^2 |M_1|^2 + \frac{1}{2}\rho Q_L^2 g_1'^2 |M_1'|^2 \right) - \frac{3}{5}g_1^2 S + \rho Q_{H_d} g_1'^2 S' \\
 \frac{dm_{\tilde{e}_R}^2}{dt} &= -8 \left(\frac{3}{5}g_1^2 |M_1|^2 + \frac{1}{2}\rho Q_E^2 g_1'^2 |M_1'|^2 \right) + \frac{6}{5}g_1^2 S + \rho Q_E g_1'^2 S', \tag{A.6}
 \end{aligned}$$

where the corresponding RGEs in the MSSM are obtained by setting $g_1' = 0$, $h_s = 0$ and $m_{\tilde{g}}^2 = 0$ everywhere. Since $U(1)'$ charges are family-universal the squared-masses of the right-handed sfermions exhibit finite splitting only by their boundary values at M_{GUT} (to the extent their Yukawa couplings can be neglected). The beta functions of the scalar mass-squareds depend on S defined in (3.7), and on

$$\begin{aligned}
 S' = \text{Tr} [m^2 Q] &= 2 \left(Q_{H_u} m_{H_u}^2 + Q_{H_d} m_{H_d}^2 + \frac{1}{2} Q_S m_S^2 \right) + \\
 & + 6Q_Q \left(m_{\tilde{t}_L}^2 + m_{\tilde{c}_L}^2 + m_{\tilde{u}_L}^2 \right) + 3Q_U \left(m_{\tilde{t}_R}^2 + m_{\tilde{c}_R}^2 + m_{\tilde{u}_R}^2 \right) + \\
 & + 3Q_D \left(m_{\tilde{b}_R}^2 + m_{\tilde{s}_R}^2 + m_{\tilde{d}_R}^2 \right) + 2Q_L \left(m_{\tilde{\tau}_L}^2 + m_{\tilde{\mu}_L}^2 + m_{\tilde{e}_L}^2 \right) + \\
 & + Q_E \left(m_{\tilde{\tau}_R}^2 + m_{\tilde{\mu}_R}^2 + m_{\tilde{e}_R}^2 \right) \tag{A.7}
 \end{aligned}$$

which vanishes if mass-squareds are universal provided that Z' -graviton-graviton anomaly cancels out i.e. $\text{Tr}[Q] = 0$.

B. Renormalization group equations in the NMSSM

In this appendix we list down RGEs for the NMSSM [20], for completeness. Since the model exhibits no gauge extension with respect to MSSM, gauge couplings and gaugino masses evolve precisely as in the MSSM (as mentioned below equations A.1 and A.4 in appendix A). On the other hand, superpotential parameters and soft masses are modified both in number and evolution, and below we provide explicit expressions for their beta functions.

The evolutions of the superpotential parameters are given by

$$\begin{aligned}
 \frac{dh_t}{dt} &= h_t \left(6h_t^2 + h_b^2 + h_s^2 - \frac{16}{3}g_3^2 - 3g_2^2 - \frac{13}{15}g_1^2 \right) \\
 \frac{dh_b}{dt} &= h_b \left(6h_b^2 + h_t^2 + h_\tau^2 + h_s^2 - \frac{16}{3}g_3^2 - 3g_2^2 - \frac{7}{15}g_1^2 \right) \\
 \frac{dh_\tau}{dt} &= h_\tau \left(4h_\tau^2 + 3h_b^2 + h_s^2 - 3g_2^2 - \frac{9}{5}g_1^2 \right) \\
 \frac{dh_s}{dt} &= h_s \left(4h_s^2 + 3h_t^2 + 3h_b^2 + h_\tau^2 + \frac{1}{2}k_s^2 - 3g_2^2 - \frac{3}{5}g_1^2 \right) \\
 \frac{dk_s}{dt} &= k_s \left(6h_s^2 + \frac{3}{2}k_s^2 \right)
 \end{aligned} \tag{B.1}$$

so that h_s and k_s exhibit a correlated RG running yet rest of the Yukawas remain as in the $U(1)'$ model (with $g_1' = 0$, of course). The MSSM limit is achieved by putting $k_s = 0$ and $h_s = 0$ while identifying $d \ln h_s/dt$ with $d \ln \mu/dt$.

The RG evolutions of the trilinear couplings are similar

$$\begin{aligned}
 \frac{dA_t}{dt} &= 2 \left(6h_t^2 A_t + h_b^2 A_b + h_s^2 A_s \right) + 2 \left(\frac{16}{3}g_3^2 M_3 + 3g_2^2 M_2 + \frac{13}{15}g_1^2 M_1 \right) \\
 \frac{dA_b}{dt} &= 2 \left(6h_b^2 A_b + h_t^2 A_t + h_\tau^2 A_\tau + h_s^2 A_s \right) + 2 \left(\frac{16}{3}g_3^2 M_3 + 3g_2^2 M_2 + \frac{7}{15}g_1^2 M_1 \right) \\
 \frac{dA_\tau}{dt} &= 2 \left(4h_\tau^2 A_\tau + 3h_b^2 A_b + h_s^2 A_s \right) + 2 \left(3g_2^2 M_2 + \frac{9}{5}g_1^2 M_1 \right) \\
 \frac{dA_s}{dt} &= 2 \left(4h_s^2 A_s + \frac{1}{2}k_s^2 A_k + 3h_t^2 A_t + 3h_b^2 A_b + h_\tau^2 A_\tau \right) + \\
 &\quad + 2 \left(3g_2^2 M_2 + \frac{3}{5}g_1^2 M_1 \right) \\
 \frac{dA_k}{dt} &= 2 \left(6h_s^2 A_s + \frac{3}{2}k_s^2 A_k \right),
 \end{aligned} \tag{B.2}$$

where the MSSM limit is obtained by putting $h_s = 0$, $k_s = 0$ everywhere and by identifying A_s with the Higgs bilinear mixing mass B .

Finally, the scalar mass-squared parameters run as follows

$$\begin{aligned}
 \frac{dm_{H_u}^2}{dt} &= 2 \left(m_{H_u}^2 + m_{H_d}^2 + m_S^2 + |A_s|^2 \right) h_s^2 + 6 \left(m_{H_u}^2 + m_{\tilde{t}_L}^2 + m_{\tilde{t}_R}^2 + |A_t|^2 \right) h_t^2 - \\
 &\quad - 8 \left(\frac{3}{4} g_2^2 |M_2|^2 + \frac{3}{20} g_1^2 |M_1|^2 \right) + \frac{3}{5} g_1^2 S \\
 \frac{dm_{H_d}^2}{dt} &= 2 \left(m_{H_u}^2 + m_{H_d}^2 + m_S^2 + |A_s|^2 \right) h_s^2 + 2 \left(m_{H_d}^2 + m_{\tilde{\tau}_L}^2 + m_{\tilde{\tau}_R}^2 + |A_\tau|^2 \right) h_\tau^2 + \\
 &\quad + 6 \left(m_{H_d}^2 + m_{\tilde{t}_L}^2 + m_{\tilde{b}_R}^2 + |A_b|^2 \right) h_b^2 - 8 \left(\frac{3}{4} g_2^2 |M_2|^2 + \frac{3}{20} g_1^2 |M_1|^2 \right) - \frac{3}{5} g_1^2 S \\
 \frac{dm_S^2}{dt} &= 4 \left(m_{H_u}^2 + m_{H_d}^2 + m_S^2 + |A_s|^2 \right) h_s^2 + \left(3m_S^2 + |A_k|^2 \right) k_s^2 \\
 \frac{dm_{\tilde{t}_L}^2}{dt} &= 2 \left(m_{\tilde{t}_L}^2 + m_{H_d}^2 + m_{\tilde{b}_R}^2 + |A_b|^2 \right) h_b^2 + 2 \left(m_{\tilde{t}_L}^2 + m_{H_u}^2 + m_{\tilde{t}_R}^2 + |A_t|^2 \right) h_t^2 - \\
 &\quad - 8 \left(\frac{4}{3} g_3^2 |M_3|^2 + \frac{3}{4} g_2^2 |M_2|^2 \right) + \frac{1}{5} g_1^2 S \\
 \frac{dm_{\tilde{t}_R}^2}{dt} &= 4 \left(m_{\tilde{t}_L}^2 + m_{H_u}^2 + m_{\tilde{t}_R}^2 + |A_t|^2 \right) h_t^2 - 8 \left(\frac{4}{3} g_3^2 |M_3|^2 + \frac{4}{15} g_1^2 |M_1|^2 \right) - \frac{4}{5} g_1^2 S \\
 \frac{dm_{\tilde{b}_R}^2}{dt} &= 4 \left(m_{\tilde{t}_L}^2 + m_{H_d}^2 + m_{\tilde{b}_R}^2 + |A_b|^2 \right) h_b^2 - 8 \left(\frac{4}{3} g_3^2 |M_3|^2 + \frac{1}{15} g_1^2 |M_1|^2 \right) + \frac{2}{5} g_1^2 S \\
 \frac{dm_{\tilde{\tau}_L}^2}{dt} &= 2 \left(m_{\tilde{\tau}_L}^2 + m_{H_d}^2 + m_{\tilde{\tau}_R}^2 + |A_\tau|^2 \right) h_\tau^2 - 8 \left(\frac{3}{4} g_2^2 |M_2|^2 + \frac{3}{20} g_1^2 |M_1|^2 \right) - \frac{3}{5} g_1^2 S \\
 \frac{dm_{\tilde{\tau}_R}^2}{dt} &= 4 \left(m_{\tilde{\tau}_L}^2 + m_{H_d}^2 + m_{\tilde{\tau}_R}^2 + |A_\tau|^2 \right) h_\tau^2 - 8 \left(\frac{3}{5} g_1^2 |M_1|^2 \right) + \frac{6}{5} g_1^2 S \\
 \frac{dm_{\tilde{u}_L}^2}{dt} &= -8 \left(\frac{4}{3} g_3^2 |M_3|^2 + \frac{3}{4} g_2^2 |M_2|^2 + \frac{1}{60} g_1^2 |M_1|^2 \right) + \frac{1}{5} g_1^2 S \\
 \frac{dm_{\tilde{u}_R}^2}{dt} &= -8 \left(\frac{4}{3} g_3^2 |M_3|^2 + \frac{4}{15} g_1^2 |M_1|^2 \right) - \frac{4}{5} g_1^2 S \\
 \frac{dm_{\tilde{d}_R}^2}{dt} &= -8 \left(\frac{4}{3} g_3^2 |M_3|^2 + \frac{1}{15} g_1^2 |M_1|^2 \right) + \frac{2}{5} g_1^2 S \\
 \frac{dm_{\tilde{e}_L}^2}{dt} &= -8 \left(\frac{3}{4} g_2^2 |M_2|^2 + \frac{3}{20} g_1^2 |M_1|^2 \right) - \frac{3}{5} g_1^2 S \\
 \frac{dm_{\tilde{e}_R}^2}{dt} &= -8 \left(\frac{3}{5} g_1^2 |M_1|^2 \right) + \frac{6}{5} g_1^2 S, \tag{B.3}
 \end{aligned}$$

where evolution of m_S^2 is modified by the singlet cubic coupling whereas rest of the squared-masses run as in the $U(1)'$ model with $g_1' = 0$.

References

- [1] S. Dimopoulos, S. Raby and F. Wilczek, *Supersymmetry and the scale of unification*, *Phys. Rev. D* **24** (1981) 1681;

- L.E. Ibáñez and G.G. Ross, *Low-energy predictions in supersymmetric grand unified theories*, *Phys. Lett.* **B 105** (1981) 439;
W.J. Marciano and G. Senjanovic, *Predictions of supersymmetric grand unified theories*, *Phys. Rev.* **D 25** (1982) 3092;
P. Langacker and N. Polonsky, *Uncertainties in coupling constant unification*, *Phys. Rev.* **D 47** (1993) 4028 [[hep-ph/9210235](#)].
- [2] L.E. Ibáñez and G.G. Ross, *SU(2)_L × U(1) symmetry breaking as a radiative effect of supersymmetry breaking in guts*, *Phys. Lett.* **B 110** (1982) 215;
K. Inoue, A. Kakuto, H. Komatsu and S. Takeshita, *Aspects of grand unified models with softly broken supersymmetry*, *Prog. Theor. Phys.* **68** (1982) 927;
L. Alvarez-Gaumé, M. Claudson and M.B. Wise, *Low-energy supersymmetry*, *Nucl. Phys.* **B 207** (1982) 96;
L. Alvarez-Gaumé, J. Polchinski and M.B. Wise, *Minimal low-energy supergravity*, *Nucl. Phys.* **B 221** (1983) 495;
J.R. Ellis, J.S. Hagelin, D.V. Nanopoulos and K. Tamvakis, *Weak symmetry breaking by radiative corrections in broken supergravity*, *Phys. Lett.* **B 125** (1983) 275;
K. Inoue, A. Kakuto, H. Komatsu and S. Takeshita, *Renormalization of supersymmetry breaking parameters revisited*, *Prog. Theor. Phys.* **71** (1984) 413.
- [3] A. Bouquet, J. Kaplan and C.A. Savoy, *On flavor mixing in broken supergravity*, *Phys. Lett.* **B 148** (1984) 69;
L.J. Hall, V.A. Kostelecky and S. Raby, *New flavor violations in supergravity models*, *Nucl. Phys.* **B 267** (1986) 415;
G.K. Leontaris, N.D. Tracas and J.D. Vergados, *Renormalization effects on flavor mixing in supergravity theories*, *Phys. Lett.* **B 206** (1988) 247.
- [4] T. Kobayashi and K. Yoshioka, *New RG-invariants of soft supersymmetry breaking parameters*, *Phys. Lett.* **B 486** (2000) 223 [[hep-ph/0004175](#)].
- [5] M. Carena, K. Huitu and T. Kobayashi, *RG-invariant sum rule in a generalization of anomaly mediated SUSY breaking models*, *Nucl. Phys.* **B 592** (2001) 164 [[hep-ph/0003187](#)];
K. Huitu, J. Laamanen and P.N. Pandita, *Sparticle spectrum and constraints in anomaly mediated supersymmetry breaking models*, *Phys. Rev.* **D 65** (2002) 115003 [[hep-ph/0203186](#)].
- [6] I. Jack, D.R.T. Jones and A. Pickering, *Renormalisation invariance and the soft β -functions*, *Phys. Lett.* **B 426** (1998) 73 [[hep-ph/9712542](#)];
N. Arkani-Hamed, G.F. Giudice, M.A. Luty and R. Rattazzi, *Supersymmetry-breaking loops from analytic continuation into superspace*, *Phys. Rev.* **D 58** (1998) 115005 [[hep-ph/9803290](#)];
I. Jack and D.R.T. Jones, *Rg invariant solutions for the soft supersymmetry breaking parameters*, *Phys. Lett.* **B 465** (1999) 148 [[hep-ph/9907255](#)];
N. Arkani-Hamed, D.E. Kaplan, H. Murayama and Y. Nomura, *Viable ultraviolet-insensitive supersymmetry breaking*, *JHEP* **02** (2001) 041 [[hep-ph/0012103](#)].
- [7] D.J. Gross and J. Wess, *Scale invariance, conformal invariance and the high-energy behavior of scattering amplitudes*, *Phys. Rev.* **D 2** (1970) 753;
J. Polchinski, *Scale and conformal invariance in quantum field theory*, *Nucl. Phys.* **B 303** (1988) 226;
A. Iorio, L. O’Raifeartaigh, I. Sachs and C. Wiesendanger, *Weyl gauging and conformal invariance*, *Nucl. Phys.* **B 495** (1997) 433 [[hep-th/9607110](#)];

- D.A. Demir, *Nonlinearly realized local scale invariance: gravity and matter*, *Phys. Lett. B* **584** (2004) 133 [[hep-ph/0401163](#)].
- [8] V.K. Dobrev and V.B. Petkova, *All positive energy unitary irreducible representations of extended conformal supersymmetry*, *Phys. Lett. B* **162** (1985) 127;
S. Minwalla, *Restrictions imposed by superconformal invariance on quantum field theories*, *Adv. Theor. Math. Phys.* **2** (1998) 781 [[hep-th/9712074](#)];
S.P. Martin and J.D. Wells, *Constraints on ultraviolet-stable fixed points in supersymmetric gauge theories*, *Phys. Rev. D* **64** (2001) 036010 [[hep-ph/0011382](#)].
- [9] A.E. Nelson and M.J. Strassler, *Suppressing flavor anarchy*, *JHEP* **09** (2000) 030 [[hep-ph/0006251](#)]; *Exact results for supersymmetric renormalization and the supersymmetric flavor problem*, *JHEP* **07** (2002) 021 [[hep-ph/0104051](#)].
- [10] P. Fayet, *Spontaneously broken supersymmetric theories of weak, electromagnetic and strong interactions*, *Phys. Lett. B* **69** (1977) 489;
J.E. Kim and H.P. Nilles, *The mu problem and the strong CP problem*, *Phys. Lett. B* **138** (1984) 150;
D. Suematsu and Y. Yamagishi, *Radiative symmetry breaking in a supersymmetric model with an extra U(1)*, *Int. J. Mod. Phys. A* **10** (1995) 4521 [[hep-ph/9411239](#)];
M. Cvetič and P. Langacker, *New gauge bosons from string models*, *Mod. Phys. Lett. A* **11** (1996) 1247 [[hep-ph/9602424](#)];
V. Jain and R. Shrock, *U(1)-a models of fermion masses without a mu problem*, [hep-ph/9507238](#);
Y. Nir, *Gauge unification, yukawa hierarchy and the μ -problem*, *Phys. Lett. B* **354** (1995) 107 [[hep-ph/9504312](#)].
- [11] J.L. Hewett and T.G. Rizzo, *Low-energy phenomenology of superstring inspired E_6 models*, *Phys. Rept.* **183** (1989) 193;
M. Cvetič and P. Langacker, *Implications of abelian extended gauge structures from string models*, *Phys. Rev. D* **54** (1996) 3570 [[hep-ph/9511378](#)];
C.T. Hill and E.H. Simmons, *Strong dynamics and electroweak symmetry breaking*, *Phys. Rept.* **381** (2003) 235 [[hep-ph/0203079](#)];
D.A. Demir and L.L. Everett, *CP-violation in supersymmetric U(1)' models*, *Phys. Rev. D* **69** (2004) 015008 [[hep-ph/0306240](#)].
- [12] M. Cvetič, D.A. Demir, J.R. Espinosa, L.L. Everett and P. Langacker, *Electroweak breaking and the mu problem in supergravity models with an additional U(1)*, *Phys. Rev. D* **56** (1997) 2861 [[hep-ph/9703317](#)].
- [13] J.R. Ellis, J.F. Gunion, H.E. Haber, L. Roszkowski and F. Zwirner, *Higgs bosons in a nonminimal supersymmetric model*, *Phys. Rev. D* **39** (1989) 844;
Y. Daikoku and D. Suematsu, *Mass bound of the lightest neutral Higgs scalar in the extra U(1) models*, *Phys. Rev. D* **62** (2000) 095006 [[hep-ph/0003205](#)];
D.A. Demir, *Distinguishing the Higgs scalars of nmssm and Z' models for large Higgs trilinear couplings*, [hep-ph/9902468](#).
- [14] G.F. Giudice and A. Masiero, *A natural solution to the mu problem in supergravity theories*, *Phys. Lett. B* **206** (1988) 480.
- [15] J.A. Casas and C. Muñoz, *A natural solution to the mu problem*, *Phys. Lett. B* **306** (1993) 288 [[hep-ph/9302227](#)].

- [16] D.A. Demir and E. Ma, *Relaxation of the dynamical gluino phase and unambiguous electric dipole moments*, *Phys. Rev. D* **62** (2000) 111901 [[hep-ph/0004148](#)];
Realistic gluino axion model consistent with supersymmetry breaking at the TeV scale, *J. Phys. G* **27** (2001) L87–L94 [[hep-ph/0101185](#)].
- [17] E.J. Chun, J.E. Kim and H.P. Nilles, *A natural solution of the μ problem with a composite axion in the hidden sector*, *Nucl. Phys. B* **370** (1992) 105.
- [18] Y. Kawamura, T. Kobayashi and J. Kubo, *Soft scalar-mass sum rule in gauge-yukawa unified models and its superstring interpretation*, *Phys. Lett. B* **405** (1997) 64 [[hep-ph/9703320](#)];
T. Kobayashi, J. Kubo, M. Mondragon and G. Zoupanos, *Constraints on finite soft supersymmetry-breaking terms*, *Nucl. Phys. B* **511** (1998) 45 [[hep-ph/9707425](#)];
D.I. Kazakov, *Finiteness of soft terms in finite $N = 1$ SUSY gauge theories*, *Phys. Lett. B* **421** (1998) 211 [[hep-ph/9709465](#)];
Y. Kawamura, T. Kobayashi and J. Kubo, *Sum rules in the superpartner spectrum of the minimal supersymmetric standard model*, *Phys. Lett. B* **432** (1998) 108 [[hep-ph/9710453](#)];
T. Kobayashi, J. Kubo and G. Zoupanos, *Further all-loop results in softly-broken supersymmetric gauge theories*, *Phys. Lett. B* **427** (1998) 291 [[hep-ph/9802267](#)].
- [19] S.P. Martin and P. Ramond, *Sparticle spectrum constraints*, *Phys. Rev. D* **48** (1993) 5365 [[hep-ph/9306314](#)];
S.P. Martin, *Generalized messengers of supersymmetry breaking and the sparticle mass spectrum*, *Phys. Rev. D* **55** (1997) 3177 [[hep-ph/9608224](#)].
- [20] J.P. Derendinger and C.A. Savoy, *Quantum effects and $SU(2) \times U(1)$ breaking in supergravity gauge theories*, *Nucl. Phys. B* **237** (1984) 307;
S.F. King and P.L. White, *Resolving the constrained minimal and next-to-minimal supersymmetric standard models*, *Phys. Rev. D* **52** (1995) 4183 [[hep-ph/9505326](#)].
- [21] G.G. Ross and R.G. Roberts, *Minimal supersymmetric unification predictions*, *Nucl. Phys. B* **377** (1992) 571;
S. Kelley, J.L. Lopez, D.V. Nanopoulos, H. Pois and K.-j. Yuan, *No scale supergravity confronts experiment*, *Phys. Lett. B* **273** (1991) 423;
D.J. Castano, E.J. Piard and P. Ramond, *Renormalization group study of the standard model and its extensions. 2. the minimal supersymmetric standard model*, *Phys. Rev. D* **49** (1994) 4882 [[hep-ph/9308335](#)];
S.P. Martin and M.T. Vaughn, *Two loop renormalization group equations for soft supersymmetry breaking couplings*, *Phys. Rev. D* **50** (1994) 2282 [[hep-ph/9311340](#)];
L.V. Avdeev, D.I. Kazakov and I.N. Kondrashuk, *Renormalizations in softly broken SUSY gauge theories*, *Nucl. Phys. B* **510** (1998) 289 [[hep-ph/9709397](#)].
- [22] D.J.H. Chung et al., *The soft supersymmetry-breaking lagrangian: theory and applications*, *Phys. Rept.* **407** (2005) 1 [[hep-ph/0312378](#)].
- [23] A.E. Nelson and L. Randall, *Naturally large $\tan \beta$* , *Phys. Lett. B* **316** (1993) 516 [[hep-ph/9308277](#)].
- [24] H. Baer, J. Ferrandis and X. Tata, *Are supersymmetric models with large $\tan \beta$ natural?*, *Phys. Lett. B* **561** (2003) 145 [[hep-ph/0211418](#)].
- [25] A.E. Faraggi, J.S. Hagelin, S. Kelley and D.V. Nanopoulos, *Sparticle spectroscopy*, *Phys. Rev. D* **45** (1992) 3272;
V.D. Barger, M.S. Berger and P. Ohmann, *The supersymmetric particle spectrum*, *Phys. Rev. D* **49** (1994) 4908 [[hep-ph/9311269](#)];

- Y. Kawamura, H. Murayama and M. Yamaguchi, *Probing symmetry breaking pattern using sfermion masses*, *Phys. Lett. B* **324** (1994) 52 [[hep-ph/9402254](#)].
- [26] J.R. Ellis, A.B. Lahanas, D.V. Nanopoulos and K. Tamvakis, *No-scale supersymmetric standard model*, *Phys. Lett. B* **134** (1984) 429;
 J.R. Ellis, C. Kounnas and D.V. Nanopoulos, *No scale supersymmetric guts*, *Nucl. Phys. B* **247** (1984) 373;
 A.B. Lahanas and D.V. Nanopoulos, *The road to no scale supergravity*, *Phys. Rept.* **145** (1987) 1.
- [27] R. Barbieri, J. Louis and M. Moretti, *Phenomenological implications of supersymmetry breaking by the dilaton*, *Phys. Lett. B* **312** (1993) 451 [[hep-ph/9305262](#)];
 A. Brignole, L.E. Ibáñez and C. Muñoz, *Towards a theory of soft terms for the supersymmetric standard model*, *Nucl. Phys. B* **422** (1994) 125 [[hep-ph/9308271](#)];
 A. Brignole, L.E. Ibáñez, C. Muñoz and C. Scheich, *Some issues in soft SUSY breaking terms from dilaton/moduli sectors*, *Z. Physik C* **74** (1997) 157 [[hep-ph/9508258](#)];
 J.A. Casas, A. Lleyda and C. Muñoz, *Problems for supersymmetry breaking by the dilaton in strings from charge and color breaking*, *Phys. Lett. B* **380** (1996) 59 [[hep-ph/9601357](#)];
 S.A. Abel, B.C. Allanach, F. Quevedo, L. Ibáñez and M. Klein, *Soft SUSY breaking, dilaton domination and intermediate scale string models*, *JHEP* **12** (2000) 026 [[hep-ph/0005260](#)].
- [28] P.G. Camara, L.E. Ibáñez and A.M. Uranga, *Flux-induced SUSY-breaking soft terms*, *Nucl. Phys. B* **689** (2004) 195 [[hep-th/0311241](#)];
 B.C. Allanach, A. Brignole and L.E. Ibáñez, *Phenomenology of a fluxed MSSM*, *JHEP* **05** (2005) 030 [[hep-ph/0502151](#)].
- [29] T. Han, P. Langacker and B. McElrath, *The Higgs sector in a $U(1)'$ extension of the MSSM*, *Phys. Rev. D* **70** (2004) 115006 [[hep-ph/0405244](#)].
- [30] J. Erler, P. Langacker and T.-j. Li, *The z - z' mass hierarchy in a supersymmetric model with a secluded $U(1)'$ -breaking sector*, *Phys. Rev. D* **66** (2002) 015002 [[hep-ph/0205001](#)].
- [31] J. Erler, *Chiral models of weak scale supersymmetry*, *Nucl. Phys. B* **586** (2000) 73 [[hep-ph/0006051](#)];
 M. Aoki and N. Oshimo, *A supersymmetric model with an extra $U(1)$ gauge symmetry*, *Phys. Rev. Lett.* **84** (2000) 5269 [[hep-ph/9907481](#)].
- [32] A.H. Chamseddine and H.K. Dreiner, *Anomaly - free gauged $U(1)'$ in local supersymmetry and baryon number violation*, *Nucl. Phys. B* **447** (1995) 195 [[hep-ph/9503454](#)].
- [33] D.A. Demir, G.L. Kane and T.T. Wang, *The minimal $U(1)'$ extension of the MSSM*, *Phys. Rev. D* **72** (2005) 015012 [[hep-ph/0503290](#)].
- [34] G.L. Kane and S.F. King, *Naturalness implications of lep results*, *Phys. Lett. B* **451** (1999) 113 [[hep-ph/9810374](#)];
 M. Bastero-Gil, C. Hugonie, S.F. King, D.P. Roy and S. Vempati, *Does lep prefer the NMSSM?*, *Phys. Lett. B* **489** (2000) 359 [[hep-ph/0006198](#)].